Research Article



The Impact of Intraperitoneal Dexmedetomidine with Bupivacaine on Patients' Postoperative Pain in Endometriosis Laparoscopic Surgery; A Randomized, Clinical Trial

Samaneh Rokhgireh¹, Abolfazl Mehdizadehkashi^{1,*}, Mansoureh Vahdat¹, Zahra Najmi², Naimeh Taghavi Shoazi¹, Vahideh Astaraii¹, Mohammad Kermansaravi³, Seyed Hamid Reza Faiz¹, Shahla Chaichian^{4,5} and Shahla Mirgaloy Bayat¹

¹Endometriosis Research Center, Iran University of Medical Sciences, Tehran, Iran ²Zanjan University of Medical Sciences, Zanjan, Iran

³Minimally Invasive Surgery Research Center, Iran University of Medical Sciences, Tehran, Iran

⁴Minimally Invasive Techniques Research Center in Women. Tehran Medical Sciences Branch. Islamic Azad University. Tehran. Iran

⁵Pars Advanced and Minimally Invasive Medical Manners Research Center, Pars Hospital, Iran University of Medical Sciences, Tehran, Iran

Corresponding author: Endometriosis Research Center, Iran University of Medical Sciences, Tehran, Iran. Email: amehdizadehkashi@yahoo.com

Received 2018 October 13; Revised 2019 January 27; Accepted 2019 February 17.

Abstract

Background: Endometriosis is routinely treated with laparoscopy, which despite significant advantages over laparotomy cannot diminish postoperative pain. Insufficient postoperative pain control decreases patient satisfaction.

Objectives: This study was designed to evaluate the efficacy of intraperitoneal dexmedetomidine (DEX) combined with bupivacaine on postoperative pain in endometriosis laparoscopic surgery.

Methods: Fifty-three patients with endometriosis, scheduled for laparoscopy in Rasoul-e-Akram Hospital, Tehran, from January 2016 to May 2017 who were randomly divided into three groups, including group 1(G1, n = 21) received 50 mL intraperitoneal saline, group 2 (G2, n = 16) received 50 mL intraperitoneal instillation of bupivacaine 0.25%, and group 3 (G3, n = 16) received 50 mL bupivacaine 0.25% plus dexmedetomidine 1 μ g/kg. Each patient with a history of allergy to local anesthetics or dexmedetomidine, cardiac disease, renal or hepatic failure, severe pulmonary disease; in addition, pregnant and comorbid obese patients were excluded from the study. Patients' postoperative pain was assessed in the recovery room after 2, 6, 12, 24, and 48 hours using visual analogue scale (VAS). Total analgesic consumption was also recorded.

Results: The postoperative VAS scores were significantly lower in group 3 than other groups in the recovery room, and 2, 6, 12, 24 and 48 hours after the surgery (P < 0.001). However, there was no significant difference between 1 and 2 groups. Furthermore, total VAS in the first 24 hours in group 3 was significantly lower than the two other groups (P < 0.001).

Conclusions: We conclude 1 μ g/kg intraperitoneal DEX administration combined with bupivacaine may prolong postoperative analgesia and decrease rescue analgesia requirement compared with bupivacaine alone.

Keywords: Bupivacaine, Dexmedetomidine, Laparoscopic Surgery, Endometriosis, Intraperitoneal Instillation

1. Background

Endometriosis is a common disease in women of reproductive age and it can present diverse symptoms, including pain and infertility (1). Despite various medical treatments, laparoscopy is the gold standard diagnostic method that provides simultaneous therapeutic management (2, 3). Although laparoscopy has great benefits compared with open procedures, postoperative pain remains an important complication (4).

Postoperative pain after laparoscopic procedures is usually a diffuse pain in abdomen, shoulder, and back

that is suggested to originate the stretching of the intraabdominal cavity (5) and the residual carbon dioxide (CO_2) in the peritoneal cavity that may cause peritoneal inflammation and phrenic nerve irritation (6). Shoulder pain is also a frequent complaint (35% to 80%) after gynecological laparoscopic surgery (7), resulting from peritoneal insufflation, mainly in exaggerated Trendelenburg position (8). Thus controlling the postoperative pain is essential, and insufficient postoperative pain control can prolong the recovery time and decrease patient satisfaction (9).

The efficacy of various analgesics, including bupiva-

Copyright © 2019, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

caine, levobupivacaine, clonidine, and naloxone have been widely investigated on postoperative pain (10). Research has revealed the ease of use and safety of local anesthetic agents that directly act on the target tissue (11, 12) and are suggested to efficiently reduce postoperative pain after laparoscopic surgery (13). Dexmedetomidine (DEX) is a α 2adrenergic agonists similar to clonidine that provides sedation, analgesia, anxiolysis, analgesia, and sympatholysis (14). The DEX has a greater α 2-receptor affinity and shorter half-life than clonidine (15) and the specificity of DEX to α 2receptors is of great interest in regional anesthesia. Moreover, the addition of DEX, as an adjuvant to bupivacaine, is proven to be effective in postoperative pain relief in various procedures such as cesarean section (16), abdominal hysterectomy (17), and other gynecologic surgeries (18, 19). However, as far as the authors are concerned, it has not been confirmed in the laparoscopic procedure for patients with endometriosis.

2. Objectives

The present study aimed to assess the analgesic property of DEX combined with bupivacaine administered intraperitoneally and compare it with the intraperitoneal administration of bupivacaine alone and normal saline in patients undergoing laparoscopy for endometriosis.

3. Methods

In the present randomized, clinical trial study, patients who were candidates of laparoscopic surgery for endometriosis in Rasoul-e-Akram Hospital, a tertiary-Medical University Center in Tehran, were recruited from January 2016 to May 2017.

After approval of the Institutional Ethics Committee (IR, IUMS, rec, 1395, 27252), STATA software was used for estimating the sample size. The test power was equivalent to 80.

$$n = \frac{\left(z_{1-}a_{/2} + z_{1-b}\right)^2 \left(s^2 + s^2\right)}{\left(\mu_1 - \mu_2\right)^2} \tag{1}$$

Fifty-three patients who met the inclusion criteria were randomly divided into three. This study was three-blinded, thus the researcher, patient, and statistical analyst were blinded to the allocation of the three groups. All patients signed the written informed consent form before the intervention.

Inclusion criteria for this study were the age between 20 - 48 years, being scheduled for elective laparoscopic surgery for the diagnosis of endometriosis, and the American Society of Anesthesiologists (ASA) class 1 and 2. Each patient with a positive history of allergy to local anesthetics or DEX, cardiac disease, renal or hepatic failure, severe pulmonary disease; in addition, pregnant and comorbid obese patients were excluded from the study. Finally, 53 patients were included in the study (Figure 1).

During operation, the subjects were monitored for vital signs using electrocardiogram, pulse-oximeter, capnography, and noninvasive blood pressure. The patients received 2 μ g/kg fentanyl + 1.5 mg midazolam intravenously (IV) and 2 mg/kg propofol + 0.2 mg/kg cisatracurium were applied to anesthesia induction. Anesthesia was maintained with 100 μ g/kg propofol. End Tidal (ET) CO₂ was maintained between 35 - 40 mmHg.

Laparoscopic procedures were performed using standard direct entry with 11-mm port at the umbilicus, while for those who required difficult deep infiltrative endometriosis interventions, 5.5 mm and 11 mm ports were placed in the bilateral lower quadrants and suprapubic zone, respectively. The patients were placed in 15 - 20°C Trendelenburg's position. During the procedure, the intraabdominal pressure was maintained at 12 - 14 mmHg. A minimum sample size of 14 was calculated to be needed in each group to achieve a study power of 80% with regard to P value less than 0.05 as a significant result. In this regard, the subjects were randomly (using randomization table) divided into three groups, including Group 1 (n = 21)received 50 mL intraperitoneal normal saline as negative control, group 2 (n = 16) received 50 mL intraperitoneal bupivacaine 0.25% as positive control, and group 3(n = 16)received 50 mL intraperitoneal bupivacaine 0.25% plus 1 μ g/kg DEX. At the end of the laparoscopic procedure, study drugs prepared in a sterile syringe were given to the researcher who was blinded to the nature and characteristics of the drugs. After homeostasis was achieved, intraperitoneal administration was guided by a camera on the port site and pelvic peritoneal surface before removal of the port in Trendelenburg's position. Then CO₂ was removed using compression of the abdomen manually at the end of the procedure. The pain severity was assessed at the recovery room, 2, 6, 12, 24, and 48 hours after the operation, by visual analogue scale (VAS) in which zero score corresponds to 'no pain' and 10 corresponds to 'maximum' or 'the worst pain'. Rescue analgesia consisted of 50 mg suppository of diclofenac sodium, administered to patients who reported VAS \geq 3 or on the patient's request. Total analgesic consumption was recorded within 48 hours.

All analyses were performed by SPSS software version 17.0 (Chicago, USA). Data were presented as mean and standard deviation (SD) for quantitative variables and number (percentage) for categorical variables. One-way analysis of variance (ANOVA) test followed by post-hoc test were used to compare quantitative variables across the groups.





Kruskal-Wallis H test was employed to compare variables with non-parametric distribution. The chi-square test was also used to determine the association between categorical variables. The P value of less than 0.05 was considered statistically significant.

4. Results

Mean age \pm SD of the patients included in the study was 32.50 \pm 7.21. There was no significant difference in terms of age and body mass index (BMI) among the three study groups (P = 0.2; Table 1).

There was one case of hysterectomy due to deep infiltrative endometriosis (DIE) in group 1, one in group 2, and two cases in group 3. There were no significant differences with respect to the type and duration of the surgery between the groups (P = 0.4; Table 2).

The postoperative VAS scores were significantly lower in group 3 than other groups in the recovery room, and 2, 6, 12, 24 and 48 hours after the surgery (P < 0.001). However, there was no significant difference between the groups of 1 and 2 (Table 1). Furthermore, total VAS in the first 24 hours was significantly lower in group 3 than the two other groups (P < 0.001). The VAS score was greater in group 2

able 1. Demographic Characteristics of the Study Population, According to the 3 Groups ^a						
	Intraperitoneal Normal Saline, Group 1 (N = 21)	Intraperitoneal Bupivacaine Group 2 (N = 16)	Intraperitoneal Bupivacaine + DEX, Group 3 (N=16)	P Value ^b		
Age, y	31 ± 6.82	32.06 ± 8.47	34.93 ± 6.06	0.2		
BMI, kg/m ²	23.87 ± 4.68	25.62 ± 3.56	26.34 ± 4.13	0.1		

Abbreviations: BMI, body mass index; DEX, Dexmedetomidine.

^aValues are expressed as mean \pm SD.

 $^{b}P < 0.05$ was considered statistically significant.

Table 2. The Frequency of Type of Procedures, According to the 3 Groups^a

	Intraperitoneal Normal Saline, Group 1 (N = 21)	Intraperitoneal Bupivacaine, Group 2 (N = 16)	Intraperitoneal Bupivacaine + DEX, Group 3 (N = 16)	P Value ^b
DIE resection	10 (47.6)	10 (62.5)	11 (68.8)	0.4
Endometrioma resection	10 (47.6)	5 (31.3)	3 (18.8)	0.4
Hysterectomy	1(4.8)	1(6.3)	2 (12.5)	0.4

^aValues are expressed as No. (%).

^bP < 0.05 was considered statistically significant.

compared with group 1; however, no significant difference (P > 0.99).

The fewest mean of total diclofenac suppositories was used in group 3 compared with the other two groups (P < 0.001) (Table 3).

We observed no significant difference in the site of maximum pain across the three groups (P = 0.3). The incidence of shoulder pain was lower in group 3 without statistically significant difference (33.3%, 50%, and 31.3% in groups 1, 2, and 3, respectively) (P > 0.05).

According to our findings, 5 patients in group 3 and 1 patient of in group 1 did not need any analgesic (Table 2).

5. Discussion

The results of the present study on the assessment of postoperative VAS scores showed significantly less pain in group 3 (DEX + bupivacaine) than group 2 (bupivacaine) and group 1 (normal saline) in the recovery room, and 2, 6, 12, 24, and 48 hours after the surgery (P < 0.001) and fewer rescue analgesic requirement in group 3. While the longest time to the first rescue analgesic requirement was in group 2. As pain relief after gynecologic laparoscopic interventions has remained a critical issue, researchers broadly investigated to determine the highest safety and tolerance to analgesic techniques among patients who planned for surgery (20). Our study demonstrated that intraperitoneally distillated DEX with bupivacaine significantly attenuated postoperative pain and declined postoperative rescue analgesic consumption compared with intraperitoneally administered bupivacaine alone or normal saline. Meta-analysis studies on minor gynecologic procegynecologic surgery has provided diverse results on the efficacy of intraperitoneal administration of bupivacaine on postoperative pain control. In this regard, Rivard et al. has found it efficient (19), while other researchers did not find any improvement in the pain control, narcotic use, length of hospital stay, or the level of patients' satisfaction following the use of bupivacaine (22, 23), which is consistent with our study findings. It may also reflect this fact that intraperitoneal normal saline and intraperitoneal bupivacaine alone have similar effectiveness, as reported by Esmat and colleagues (6). As hypothesized, the local anesthetic agent administration causes visceral afferent signaling, and modification of visceral nociception that will block sodium channels (24). Thus recent studies have evaluated the efficacy of the combination of intraperitoneal bupivacaine with another local anesthetic agent. Ahmed et al. showed that intraperitoneal instillation of meperidine or DEX in combination with bupivacaine 0.25% significantly decreased the postoperative analgesic requirements and the incidence of shoulder pain compared with bupivacaine 0.25% alone in patients undergoing laparoscopic gynecological surgeries (25). Memis et al. also demonstrated that administrating a combination of clonidine and bupivacaine via intra-peritoneal root could provide more effective analgesia compared to bupivacaine alone during the early postoperative period (26). Narasimham and Rao also found that the intraperitoneal administration of bupivacaine alone or combined with DEX or tramadol could relieve procedure-related fol-

dures (such as tubal ligation) confirmed the efficacy of in-

traperitoneal bupivacaine on postoperative pain (21). How-

ever, research on patients undergoing minimally invasive

Table 3. Total Analgesic Consumption in 48 Hours Postoperatively								
	Intraperitoneal Normal Saline, Group 1 (N = 21)	Intraperitoneal Bupivacaine, Group 2 (N = 16)	Intraperitoneal Bupivacaine +DEX, Group 3 (N = 16)	P Value				
Diclofenac consumption	2±1	3.8 ± 0.98	1.18 ± 1.16	< 0.0001				

lowing after laparoscopic cholecystectomy (27). The prolonged sedation of DEX is attributed to its longer half-life than clonidine (28). The results of the above-mentioned studies are consistent with the results of the present study, confirming the efficacy of the combination of DEX with another analgesic; however, the surgical type and the adjuvant analgesic differ among studies.

Similar to the results of the present study, other researchers have also revealed that intraperitoneal administration of DEX with bupivacaine, in laparoscopic gynecologic surgeries, was associated with a reduction in VAS and postoperative analgesic requirements in the hours of surgery (29-31). In addition to these results, the present study showed maximum painlessness at 48th hour postoperatively in DEX plus bupivacaine group.

Additionally, there was no statistically significant difference in the place of maximum pain in 24 hours and administration of local anesthetic had no efficiency on trocar site pain. As far as the authors are concerned, the combination of DEX with bupivacaine has not been studied on the laparoscopic procedure for the treatment of endometriosis and the origin of postoperative pain depends on the surgical technique. Therefore, further studies on the pain origin of this specific type of surgery may elucidate the mode of analgesia in the present study.

The strengthening point of the present study was comprised of a combination of gynecologic procedures such as retroperitoneal dissection, endometrioma, endometriotic nodule resection, and hysterectomy; all patients were postoperatively planned for similar pain regimen, suggesting the differences in pain controlling due to administrating intra-peritoneal local anesthetics. Also, the extended follow-up period of 48 hours (compared to 24 hours in previous studies) is the other strength of our study. The limitation of our study included small sample size and all surgeries were not performed by a single surgeon; however, the surgical techniques and equipment were unified.

As there are few studies on using a combination of DEX and bupivacaine, further studies considering different dosages and roots of local anesthetics are required to compare this approach of analgesia with other regional techniques with the goal of achieving more benefit with regard to postoperative pain relief along with minimized adverse effects following laparoscopic surgeries.

5.1. Conclusions

Our study showed that the administration of the intraperitoneal local anesthetic agent, DEX, in combination with bupivacaine can effectively reduce the need for prescribing postoperative analgesic drugs as well as decrease postoperative pain severity in minimally invasive laparoscopic surgery.

Acknowledgments

The authors sincerely thanks to Iran University of Medical Sciences, Tehran, Iran.

Footnotes

Authors' Contribution: Samaneh Rokhgireh designed the study. Abolfazl Mehdizadehkashi supervised the project. Mansoureh Vahdat, Zahra Najmi, Naimeh Taghavi Shoazi, and Vahideh Astaraii helped to data gathering. Mohammad Kermansaravi and Seyed Hamid Reza Faiz analyzed data. Shahla Chaichian and Shahla Mirgaloy Bayat prepared the manuscript.

Clinical Trial Registration: IRCT20150817023666N11.

Conflict of Interests: The authors have no conflict of interest.

Ethical Considerations: The authors have respected to ethical and legal considerations. Ethical code: IR, IUMS, rec, 1395, 27252.

Funding/Support: This study was financially supported by Iran University of Medical Sciences.

References

- McLeod BS, Retzloff MG. Epidemiology of endometriosis: An assessment of risk factors. *Clin Obstet Gynecol.* 2010;**53**(2):389–96. doi: 10.1097/GRF.0b013e3181db7bde. [PubMed: 20436315].
- Duffy JM, Arambage K, Correa FJ, Olive D, Garry R, Barlow DH, et al. Laparoscopic surgery for endometriosis. *Cochrane Database Syst Rev.* 2014;4. doi: 10.1002/14651858.cd011031.
- Singh SS, Suen MW. Surgery for endometriosis: Beyond medical therapies. *Fertil Steril.* 2017;107(3):549–54. doi: 10.1016/j.fertnstert.2017.01.001. [PubMed: 28189295].
- Schneider A, Altgassen C, Hertel H, Tozzi R, Koehler C. Advantages of laparoscopy. Int J Gynecol Cancer. 2003;13(Suppl 1):121.2–21. doi: 10.1136/ijgc-00009577-200303001-00452.

- Hernandez-Palazon J, Tortosa JA, Nuno de la Rosa V, Gimenez-Viudes J, Ramirez G, Robles R. Intraperitoneal application of bupivacaine plus morphine for pain relief after laparoscopic cholecystectomy. *Eur J Anaesthesiol.* 2003;20(11):891–6. doi: 10.1017/S0265021503001431. [PubMed: 14649341].
- Esmat ME, Elsebae MM, Nasr MM, Elsebaie SB. Combined low pressure pneumoperitoneum and intraperitoneal infusion of normal saline for reducing shoulder tip pain following laparoscopic cholecystectomy. *World J Surg.* 2006;**30**(11):1969–73. doi: 10.1007/s00268-005-0752z. [PubMed: 17043939].
- Kaloo P, Hills R, Kaloo C, Whittaker M. Interventions for the reduction of shoulder pain following gynaecological laparoscopic procedures. *Cochrane Database Syst Rev.* 2014. doi: 10.1002/14651858.cd011101.
- Jabbour-Khoury SI, Dabbous AS, Gerges FJ, Azar MS, Ayoub CM, Khoury GS. Intraperitoneal and intravenous routes for pain relief in laparoscopic cholecystectomy. *JSLS*. 2005;9(3):316–21. [PubMed: 16121879]. [PubMed Central: PMC3015602].
- Kim NS, Lee JS, Park SY, Ryu A, Chun HR, Chung HS, et al. Oxycodone versus fentanyl for intravenous patient-controlled analgesia after laparoscopic supracervical hysterectomy: A prospective, randomized, double-blind study. *Medicine (Baltimore)*. 2017;96(10). e6286. doi:10.1097/MD.00000000006286. [PubMed: 28272250]. [PubMed Central: PMC5348198].
- Abdulla S, Eckhardt R, Netter U, Abdulla W. A randomized, doubleblind, controlled trial on non-opioid analgesics and opioid consumption for postoperative pain relief after laparoscopic cholecystectomy. *Acta anaesth Belg.* 2012;63(1):43–50.
- Salihoglu Z, Yildirim M, Demiroluk S, Kaya G, Karatas A, Ertem M, et al. Evaluation of intravenous paracetamol administration on postoperative pain and recovery characteristics in patients undergoing laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech.* 2009;19(4):321–3. doi: 10.1097/SLE.0b013e3181b13933. [PubMed: 19692881].
- Kehlet H, Liu SS. Continuous local anesthetic wound infusion to improve postoperative outcome: back to the periphery? *Anesthesiology*. 2007;**107**(3):369–71. doi: 10.1097/01.anes.0000278901.96504.49. [PubMed: 17721238].
- Golubović S, Golubović V, Cindrić-Stančin M, Sotošek Tokmadžić V. Intraperitoneal analgesia for laparoscopic cholecystectomy: Bupivacaine versus bupivacaine with tramadol. *Collegium Antropologicum*. 2009;**33**(1):299–302.
- Brummett CM, Wagner DS. The use of alpha-2 agonists in peripheral nerve blocks: A review of the history of clonidine and a look at a possible future for dexmedetomidine. *Semin Anesth Perioperative Med Pain*. 2006;25(2):84–92. doi: 10.1053/j.sane.2006.02.001.
- Savola JM, Ruskoaho H, Puurunen J, Salonen JS, Karki NT. Evidence for medetomidine as a selective and potent agonist at alpha 2-adrenoreceptors. J Auton Pharmacol. 1986;6(4):275–84. doi: 10.1111/j.1474-8673.1986.tb00654.x. [PubMed: 2880852].
- Li Z, Tian M, Zhang CY, Li AZ, Huang AJ, Shi CX, et al. A randomised controlled trial to evaluate the effectiveness of intrathecal bupivacaine combined with different adjuvants (fentanyl, clonidine and dexmedetomidine) in caesarean section. *Drug Res (Stuttg)*. 2015;65(11):581–6. doi: 10.1055/s-0034-1395614. [PubMed: 25504002].
- Almarakbi WA, Kaki AM. Addition of dexmedetomidine to bupivacaine in transversus abdominis plane block potentiates post-operative pain relief among abdominal hysterectomy patients: A prospective randomized controlled trial. *Saudi J Anaesth.* 2014;8(2):161–6. doi: 10.4103/1658-354X.130683. [PubMed: 24843325]. [PubMed Central: PMC4024669].
- Ahmed B, Abd Elmawgoud MA, Doaa R. Antinociceptive effect of (a2adrenoceptor agonist) dexmedetomidine vs meperidine, topically,

after laparoscopic gynecologic surgery. J Med Sci. 2008;8(4):400-4. doi:10.3923/jms.2008.400.404.

- Rivard C, Vogel RI, Teoh D. Effect of intraperitoneal bupivacaine on postoperative pain in the gynecologic oncology patient. *J Minim Invasive Gynecol.* 2015;22(7):1260–5. doi: 10.1016/j.jmig.2015.07.013. [PubMed: 26216095]. [PubMed Central: PMC4631626].
- Jimenez Cruz J, Diebolder H, Dogan A, Mothes A, Rengsberger M, Hartmann M, et al. Combination of pre-emptive port-site and intraoperative intraperitoneal ropivacaine for reduction of postoperative pain: A prospective cohort study. *Eur J Obstet Gynecol Reprod Biol*. 2014;**179**:11–6. doi: 10.1016/j.ejogrb.2014.05.001. [PubMed: 24965972].
- Harrison MS, DiNapoli MN, Westhoff CL. Reducing postoperative pain after tubal ligation with rings or clips: A systematic review and meta-analysis. *Obstet Gynecol.* 2014;**124**(1):68–75. doi: 10.1097/AOG.0000000000342. [PubMed: 24901271].
- Arden D, Seifert E, Donnellan N, Guido R, Lee T, Mansuria S. Intraperitoneal instillation of bupivacaine for reduction of postoperative pain after laparoscopic hysterectomy: A double-blind randomized controlled trial. *J Minim Invasive Gynecol*. 2013;20(5):620–6. doi: 10.1016/j.jmig.2013.03.012. [PubMed: 23706546].
- Zmora O, Stolik-Dollberg O, Bar-Zakai B, Rosin D, Kuriansky J, Shabtai M, et al. Intraperitoneal bupivacaine does not attenuate pain following laparoscopic cholecystectomy. *JSLS*. 2000;4(4):301-4. [PubMed: 11051189].
- Kahokehr A, Sammour T, Srinivasa S, Hill AG. Systematic review and meta-analysis of intraperitoneal local anaesthetic for pain reduction after laparoscopic gastric procedures. *Br J Surg.* 2011;**98**(1):29–36. doi: 10.1002/bjs.7293. [PubMed: 20979101].
- Shukla U, Prabhakar T, Malhotra K, Srivastava D, Malhotra K. Intraperitoneal bupivacaine alone or with dexmedetomidine or tramadol for post-operative analgesia following laparoscopic cholecystectomy: A comparative evaluation. *Indian J Anaesth*. 2015;**59**(4):234– 9. doi: 10.4103/0019-5049.155001. [PubMed: 25937650]. [PubMed Central: PMC4408652].
- Memis D, Turan A, Karamanlioglu B, Tukenmez B, Pamukcu Z. The effect of tramadol or clonidine added to intraperitoneal bupivacaine on postoperative pain in total abdominal hysterectomy. *J Opioid Manag.* 2005;1(2):77–82. doi: 10.5055/jom.2005.0020. [PubMed: 17319251].
- Narasimham ML, Rao BD. Comparison of intraperitoneal bupivacaine alone or with dexmedetomidine or tramadol for post-operative analgesia following laparoscopic cholecystectomy. *Indian J Clin Anaesth.* 2017;4(1):143–6. doi: 10.18231/2394-4994.2017.0030.
- Virtanen R, Savola JM, Saano V, Nyman L. Characterization of the selectivity, specificity and potency of medetomidine as an alpha 2-adrenoceptor agonist. *Eur J Pharmacol.* 1988;**150**(1-2):9-14. doi: 10.1016/0014-2999(88)90744-3. [PubMed: 2900154].
- Oza VP, Parmar V, Badheka J, Nanavati DS, Taur P, Rajyaguru AM. Comparative study of postoperative analgesic effect of intraperitoneal instillation of dexmedetomidine with bupivacaine and bupivacaine alone after laparoscopic surgery. *J Minim Access Surg.* 2016;**12**(3):260–4. doi: 10.4103/0972-9941.181370. [PubMed: 27279399]. [PubMed Central: PMC4916754].
- Fares KM, Mohamed SA, Abd El-Rahman AM, Mohamed AA, Amin AT. Efficacy and safety of intraperitoneal dexmedetomidine with bupivacaine in laparoscopic colorectal cancer surgery, a randomized trial. *Pain Med.* 2015;**16**(6):1186–94. doi: 10.1111/pme.12687. [PubMed: 25585502].
- Pati BK. Intraperitoneal analgesia for postoperative pain relief after laparoscopic gynecological surgeries. Int J Reprod Contracept Obstet Gynecol. 2017;6(11):5099. doi: 10.18203/2320-1770.ijrcog20175032.