



The Prevalence of Depression and Its Association with Self-Management Behaviors in Chronic Disease Patients

Seyde Shahrbanoo Daniali¹, Firoozeh Mostafavi Darani¹, Elahe Tavassoli², Atefe Afshari¹, Farideh Forouzande³ and Ahmad Ali Eslami^{1,*}

¹Department of Health Education & Promotion, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran

²Department of Public Health, School of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

³Isfahan Health Care Center, Isfahan, Iran

*Corresponding author: Department of Health Education & Promotion, Isfahan University of Medical Sciences, Isfahan, Iran. Tel: +98-9132118364, Fax: +98-3136688628, E-mail: eslamiaa@gmail.com

Received 2016 December 24; Revised 2018 March 02; Accepted 2018 October 05.

Abstract

Background: Co-morbid depression in patients with chronic disease is associated with weak self-care and high complications.

Objective: The study aimed to determine the relationship of self-care behaviors and the kind of chronic disease with depression in adults with chronic disease in Isfahan, Iran.

Methods: This cross-sectional study was conducted in government healthcare centers. A total of 483 eligible consent respondents participated from March 2015 to June 2015. The “Stanford self-management questionnaire” and the patient health questionnaire (PHQ-9) were used for the assessment of self-management behaviors and depression. Data were analyzed using the SPSS 17 software.

Results: The prevalence of depression in participants was 14.7%. There was a significant relationship between self-management behaviors such as physical activity ($P < 0.05$) and medication adherence ($P < 0.001$), and depression. The risk of depression was greater in rheumatoid arthritis disease patients (OR= 4.41, 95% CI (0.47 - 5.38), $P > 0.001$) and skeletal disease patients (OR= 1.70, 95% CI (1.03 - 2.77), $P < 0.05$) than in other chronic disease patients.

Conclusions: We found a negative association between depression and self-care activities in Iranian patients with chronic diseases. Psychological treatment for depression should be considered to improve the prognosis of chronic diseases.

Keywords: Depression, Chronic Disease, Medication Adherence, Exercise, Nutritional Status, Self-Management

1. Background

Depression will account for the second largest cause of disease burden by 2020 (1). In addition, two-thirds of all deaths, disability-adjusted life years, and the burden on the national health care system are mostly linked to chronic diseases worldwide (2). Obviously, chronic diseases and their treatment affect the patient’s physical and mental health so that the evaluated risk of depression is reported between 8.3% among diabetic and 44% in kidney disease patients (3, 4). In some studies, symptoms of depression were reported twice more than the adults in the general population (5). These patients have a lower health status and more complications, mortality, and medical costs, too (6).

Since the chronic disease has a long period and incomplete cure, patients should be able to manage their disease to have a good quality of life. The patients with chronic illness should be considered on self-management. In fact,

the outcome of chronic illness depends on effective self-management (7).

Self-management means “the tasks of having confidence in patients to deal with medical management, role management, and emotional management of their conditions” (8).

On the other hand, psychological factors can affect self-management behaviors such as medication adherence (9), physical activity (10), medical appointments (11), and help-seeking (12). In addition, clinician-patient communication is restricted in the presence of depression in these patients (13).

To the best of our knowledge, in spite of increasing chronic diseases in our country, the prevalence of depression and its association with chronic diseases have been discussed a little. In addition, available studies have focused on depression only among diabetic (14), renal (15, 16), or heart disease (17) patients separately. However, so far,

very little attention has been paid to patients with various types of chronic diseases simultaneously.

This paper seeks to address the following questions: (1) the effect of the severity of depression on self-care behaviors (physical activity, medication adherence) and self-management (communication with the physician or other health care professionals, and daily activities among the outpatients), (2) the association between demographic and contextual variables (age, sex, duration of illness, type of disease, marital, and economic status) and depression, and (3) the differences among the severity of physical symptoms such as pain in the presence of depression in the government healthcare centers in the urban areas of Isfahan, Iran.

2. Objective

The aim of the present study is to explain the relationship between severity of depression and self-care behaviors in Iranian patients with chronic diseases.

3. Materials and Methods

3.1. Subjects

The present study was a cross-sectional study. The statistical population of this study included patients with any kind of chronic diseases admitted to the government healthcare centers of urban areas of Isfahan, Iran. These patients included middle-aged individuals or elders that registered as chronic disease patients in these centers. Based on the national care protocols, their health history was recorded in the health records and the presence of the disease was approved by the physicians of the health care centers to receive care and treatment. According to the Cochran formula by considering the respective levels of confidence interval and probability (0.95 and 0.5), the error level of 0.04, and the power statistical test of 80%, the sample size was calculated as 496. The sample was selected based on a two-step method. First, six healthcare centers were selected randomly. Since the number of registered patients in each healthcare center was different, 25 - 50 patients having the inclusion criteria were selected randomly to participate in the study from every healthcare center. The inclusion criteria were having a chronic illness at least for six months ago, having the prescription of medication, willingness to take part in the survey, and having the ability to walk. The patients with contradictory answers due to geriatric problems and distorted questionnaires were excluded so that 13 patients were omitted from the study.

After obtaining an ethical approval from the ethics committee of Isfahan University of Medical Sciences (reference: 393790.12/1/2014), coordinating with the related units, and obtaining necessary permissions, informed consent forms were filled out by the patients. They were assured that the questionnaires were anonymous so that the number and the type of disease would be held confidential. Questionnaires were completed during interviews in a quiet room (the majority of patients had not adequate ability to fill out the questionnaires because of illiteracy or vision problems). To avoid bias, one trained questioner conducted all of the interviews.

3.2. Study Tool

We measured social and demographic data with a set of questions related to age, marital status, sex, education level, weight, height, type of disease (diabetes, hypertension, heart disease, osteoporosis, arthritis, cancer, hypothyroidism, renal disease, and so on), and duration of illness. Their weight was recorded in kilograms using a calibrated digital scale. Participants were weighed without their shoes. Their heights were obtained by taking the average of two readings of height in meters using a portable stadiometer. Body mass index was determined by dividing weight (kg) by height squared (m^2).

The translated version of "Stanford Self-Management questionnaire in chronic disease patients" was used. This instrument was developed and validated by the Stanford Patient Education Research Center. The questionnaire evaluates concepts such as self-efficacy, communication with health care workers, social/daily role activity limitation, physical symptoms, and other self-care activities related to chronic illness (18). We followed the translation-back-translation process. Then, nine experts were invited to assess the face and content validity indices. The Persian version of the questionnaire was revised according to their comments. After the pilot test, the reliability coefficient of the total scale was $\alpha = 0.79$ with the internal reliability of 0.76. The questionnaires were filled out by 30 patients with at least one chronic disease. The reliability coefficient of the total scale was $\alpha = 0.79$; then, these patients were omitted from the main study (19).

Depression was evaluated by the nine-item Patient Health Questionnaire (PHQ-9) as an easy-to-use and acceptable instrument for depression measurement. The questionnaire scores each of the nine DSM-IV criteria for depression as "0" (not at all) to "3" (nearly every day). PHQ-9 scores of 5, 10, 15, and 20 represent mild, moderate, moderately severe, and severe depression, respectively (20). Internal reliability has been reported with Cronbach's alpha of 0.89 (18).

The Visual Analogue Scale for Pain is a numeric scale to evaluate physical symptoms (pain) intensity and its internal consistency was 0.86 among patients with a kind of chronic illness (21).

3.3. Social/Daily Role Activity Limitation

Four questions measured the effects of illness on daily activities such as social relationships with family, neighbors, friends, and other impacts of the illness on hobbies, the indoors duties and outside house tasks during the two past weeks. The obtained internal correlation for the scales varied from 0.70 - 0.75 (18).

The respondents were asked to answer several questions about patient self-care behaviors, including:

1. Physical activity: Did you participate in walking or other aerobic exercises as an exercise during the two past weeks (not as leisure time activity). The choices included never, < 30, 30 - 60, 60 - 180, and more than 180 minutes with the respective scores of 0 - 4 (19).

2. Healthy eating was assessed by the question regarding the consumption of 5 fruits and vegetable servings and eating fat or fried meals daily that scored as "0" (not at all) to "5" (always) (19).

3. Medication Adherence was measured by a 4-item questionnaire. These questions were related to forgiveness and suspicion of drug use discontinuation due to healing or worsening of general health during the two past weeks. The items scoring was based on yes (1) or no (0) (21).

Data were analyzed using Statistical Package for the Social Sciences (SPSS; version 17, SPSS Inc., Chicago, IL). Descriptive statistics were used to establish the frequency, range, mean, and standard deviation (SD) of demographics and physical symptoms (pain, stress) in depression and non-depression groups. The Chi-square test was used to evaluate differences in qualitative variables (gender, economic status, marital status, age, and education level) and self-management behaviors (physical activity, medication adherence, healthy nutrition, and communication with physicians (or health care professionals) in terms of depression. Spearman correlation was undertaken to examine relationships between depression, demographics, self-care duties, and social/daily role activity.

4. Results

From among eligible patients approached, 483 subjects were successfully recruited with a response rate of 97%. Three percent were excluded from the study due to the questionnaire's distortion. The age range of participants was 30 - 76 years, with a mean of 55.15 (7.21) years and more

than half of them were older than 50 years. Nearly 90% of the respondents were overweight or obese. The duration of disease was between 6 months and 45 years with a mean of 8 years for 1 - 5 chronic illnesses. Two-thirds of the participants had approximately more than one chronic disease and their economic status was moderate. Other demographic characteristics are presented in Table 1.

Based on the self-report questionnaire, the majority of participants (85.2%) were not depressed and 14.7% were depressed. Twenty six cases (5.4%) had moderately severe or severe depression. Majority of the depressed participants were female and only 8.3% of them were male ($P < 0.05$). The mean age of the patients with depression was significantly lower than that of the non-depressed patients ($P < 0.05$).

The Chi-square test showed that there was a significant relationship between self-care behaviors such as physical activity ($P < 0.05$) and medication adherence ($P = 0.001$), and depression. In addition, the relationship between depression and social role activity was significant ($P < 0.001$). There was a relationship between depression and self-care activities so that the more severity of depression, the less physical activity and medication adherence. There was a significant relationship between other behaviors such as social/daily role activities and depression, too. Nearly 20% of the patients had a weak medication adherence because of forgiveness or doubt to take medication. In addition, disruption in social role activity increased with the severity of depression (Figure 1). There was no significant relationship between healthy eating and depression (Table 2). In our study, one-fifth of these patients had problems in communication with family, relatives, and doing their household chores.

There was a significant relationship between the presence of diseases such as rheumatoid arthritis or other skeletal diseases (osteoporosis (and depression. Rheumatoid arthritis patients were more likely to report depression (OR = 4.41, 95% CI 0.47 - 1.38, $P < 0.05$) that was the case of skeletal disease patients (OR = 1.70, 95%CI 1.03 - 2.77, ($P < 0.05$). In other words, the risk of depression in rheumatoid arthritis and skeletal disease patients was 4.41 and 1.70, respectively, which was greater than the risk of depression in other chronic disease patients. Overall, the level of pain was greater among depressive patients than among non-depressive patients (Table 3).

There was a significant relationship between economic status and depression ($P < 0.05$), so that greater percentage of patients with an unfavorable economic level reported a higher level of depression ($P < 0.05$) (Table 1).

Table 1. Demographic Characteristics of Depressive and Non-Depressive Outpatients in Healthcare Centers (N = 483)

Factors	Total (N = 483) ^a	Non-Depression (N = 404) ^a	Depression (N = 79) ^a	P Value ^b
Sex				0.003 ^b
Female	412 (85.3)	336 (83.2)	76 (96.2)	
Male	71 (14.7)	68 (16.8)	3 (3.8)	
Age, y (mean ± SD)	55.15 ± 7.21	55.53 ± 7.02	55.28 ± 7.92	0.5
BMI (mean ± SD)	29.71 ± 4.37	29.59 ± 4.28	30.39 ± 4.78	0.136
BMI ranges				0.553
18.5 - 24.99	51 (10.6)	43 (10.6)	8 (10.1)	
25 - 29.99	226 (46.8)	193 (47.8)	33 (41.8)	
< 30	206 (42.7)	168 (41.6)	38 (48.1)	
Self-evaluation of economic status				0.005 ^b
Bad	95 (19.7)	69 (17.1)	26 (32.9)	
Moderate	350 (72.5)	301 (74.5)	49 (62.0)	
Good	38 (7.9)	34 (8.4)	4 (5.1)	
Education status				0.219
Illiterate	176 (36.4)	144 (35.6)	32 (40.5)	
Under 12 years of education	233 (48.2)	192 (47.5)	41 (51.9)	
12 years of education	58 (12)	53 (13.1)	5 (6.3)	
Academic education	16 (3.3)	15 (3.7)	1 (1.3)	
Marital status				0.7
Married	432 (89.4)	362 (89.6)	70 (88.6)	
Divorced or widow	79 (16.4)	70 (17.6)	9 (16.2)	

^aValues are expressed as No. (%) unless otherwise indicated.

^bThe significance level of Comparison between the two groups of depression and non-depression.

Table 2. The Association Between Depression Status and Self-Management in Terms of Depression Among Outpatients in Healthcare Centers (N = 483)

Self-Care Activities (Past Two Weeks)	Non-Depression (N = 404) ^a	Depression (N = 79) ^a	OR (95% CI)	P Value
Five servings of fruits and vegetables daily	395 (97.8)	78 (98.7)	1.77 (0.22 - 14.23)	0.583
Eating fat or fried meals	177 (43.8)	27 (34.2)	0.67 (0.40 - 1.10)	0.113
Physical activity (30 - 60 min/day)	75 (18.6)	6 (7.6)	0.36 (0.15 - 0.86)	0.017
Medication adherence	275 (68.1)	39 (49.4)	0.46 (0.28 - 0.74)	0.001
Disruption in daily activities and social communication	33 (8.2)	31 (39.2)	7.26 (4.08 - 12.90)	< 0.001

Abbreviation: CI, confidence interval.

^aValues are expressed as No. (%).

5. Discussion

This study was conducted among 489 outpatients in government health care centers of urban areas of Isfahan, Iran. In our study, there was a significant relationship between depression and self-management behaviors (medication adherence, physical activity, social/daily role activity) among chronic disease patients. The remarkable point is that increased depression severity is followed by weak self-care behaviors such as physical activity. Similar results have been reported by other researchers (22-24). Depression is also known as a mediator for medication adherence in diabetic and hypertensive patients (25, 26). It seems the nature of prolonged diseases, higher perceived barriers,

the outlook of hopelessness, and lack of energy and concentration among chronic disease patients can affect medication adherence. Therefore, medication adherence is low in these patients.

There was no association between healthy nutrition and depression in the current study. However, the relationship between depression and a dietary pattern containing vegetables, fruits, and the low-fat food was established in non-patients (25, 26) so that some studies pointed to the protective role of vegetables against depressive symptoms (27). Nevertheless, Rivera-Hernandez's study was in line with our study (28). This difference may be related to the type of questionnaire used while Jacka et al. (27) investigated the dietary pattern with an accurate instrument of

Table 3. The Association Between Characteristics of Chronic Diseases and Depression Status Among Outpatients in Healthcare Centers (N = 483)


Factors	Total (N = 483) ^a	Non-Depression (N = 404) ^a	Depression (N = 79) ^a	OR (95% CI)	P Value ^b
Name of disease^c					
Hypertension	359 (74.3)	303 (75.0)	56 (70.9)	0.81 (0.47-1.38)	0.444
Diabetes	283 (56.6)	239 (59.2)	44 (55.7)	0.87 (0.53-1.41)	0.568
Osteoporosis	144 (29.8)	115 (28.5)	29 (36.7)	1.46 (0.88-2.42)	0.143
Heart disease	124 (25.7)	98 (24.3)	26 (32.9)	1.53 (0.91-2.58)	0.107
Renal disease	73 (15.1)	58 (14.4)	15 (19.0)	1.39 (0.75-1.62)	0.293
Arthritis rheumatoid	36 (7.9)	22 (5.4)	16 (20.3)	4.41 (2.19-8.85)	< 0.001
Skeletal disease	164 (34.0)	129 (31.9)	35 (44.3)	1.70 (1.03-2.77)	0.034
Duration disease, y	8.06 ± 6.36	8.01 ± 6.23	8.32 ± 7.05	1.01 (0.97-1.05)	0.697
Pain (0-10)	4.93 ± 3.69	4.46 ± 3.64	7.34 ± 3.01	1.27 (1.17-1.38)	< 0.001

Abbreviation: CI, confidence interval.

^aValues are expressed as No. (%) or mean ± SD.

^bThe significance level of comparison between the two groups of depression and non-depression.

^cSome of patients had two or more diseases.

**Figure 1.** The mean score of depression, some self-management behaviors, and social/role activity among the outpatients in health care centers (N = 483).

more than 100 food items (28). On the other hand, careful consideration of nutritional patterns among patients is required to respond to this difference.

In our study, there was a relationship between pain and other painful diseases such as rheumatoid arthritis and other skeletal diseases. Studies have found that patients with more frequent pain episodes or longer pain duration are several times more likely to be depressed compared to other patients (29). Major depressive patients had a higher level of pain than non-depressive patients (29). In studies, depression was known as a strong predictor of the onset of an episode of intense low back pain (30). As the history of

depression was not discussed in our study, we cannot exactly determine whether pain leads to depression or vice versa.

Severe depression was seen in 5.4% of the participants and major depression in 8.3% by Li (3). Kaur et al. pointed to a similar prevalence among diabetic patients (31). This is while the higher rates of depression have been reported by Sharif et al. (16) and Abbas Tavallai et al. (15). This difference may be attributed to different instruments and sample sizes that included hospitalized patients with heart or renal diseases. In the current study, the majority of participants were diabetic or hypertensive outpatients without

serious complications. In the present study, one-fifth of the participants complained especially of the limitations that prevented from the achievement of household tasks and restrictions in social relationships. Other studies are in line with our study so that the presence of depression in hypertensive patients even without depression symptoms was followed by a higher level of disability (32). In fact, the restriction of social role-related activities is attributed to negative feelings induced by the nature of the long-term disease. The women and patients with low economic status were more depressed than the others in this study. Other studies revealed that sex, low socio-economic class, and marital status were the predictors of depression, too (3, 32, 33).

Definitely, adherence to treatment advice in chronic diseases can decrease the mortality and morbidity of the disease. On the other hand, the influence of psychological factors is emphasized in a better self-care.

5.1. Restriction of Study

Sampling from healthcare centers may be thought to limit the generality of the findings. We had no prior history of depression or antidepressant consumption in patients, too.

5.2. Conclusions

We found a positive association between depression and self-care activities in Iranian patients with chronic diseases. Psychological treatment for depression should be considered to improve the prognosis of chronic diseases.

Acknowledgments

We hereby express our gratitude to the Research Deputy of Isfahan University of Medical Sciences, Department of Public Health, and all of the personnel of public health centers. We also express our thanks to all the patients who participated in our study and to other expert committees for their collaboration.

Footnotes

Authors' Contribution: Seyde Shahrbanoo Daniali and Elahe Tavassoli developed the design, acquisition of data, analysis of data and drafting the article, original idea, statistical analysis, and manuscript revision. Atefe Afshari performed drafting the article and significant revisions. Firoozeh Mostafavi Darani performed analysis and interpretation of data. Ahmad Ali Eslami contributed final approval of the version to be published, to be accountable for

all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Clinical Trial Registration Code: None Declared.

Declaration of Interest: The authors declare that they have no conflict of interest.

Funding/Support: We don't receive any financial support for publication.

Ethical Considerations: The ethical approval of ethical committee (reference: 393790.12/1/2014) was taken from Isfahan University of Medical Sciences.

References

- Murray CJL, Lopez AD; Harvard School of Public Health; World Health Organization, World Bank. *The global burden of disease: A comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020*. Cambridge, MA: Harvard University Press; 1996.
- Palmer S, Vecchio M, Craig JC, Tonelli M, Johnson DW, Nicolucci A, et al. Prevalence of depression in chronic kidney disease: Systematic review and meta-analysis of observational studies. *Kidney Int*. 2013;**84**(1):179–91. doi: [10.1038/ki.2013.77](https://doi.org/10.1038/ki.2013.77). [PubMed: [23486521](https://pubmed.ncbi.nlm.nih.gov/23486521/)].
- Li C, Ford ES, Strine TW, Mokdad AH. Prevalence of depression among U.S. adults with diabetes: Findings from the 2006 behavioral risk factor surveillance system. *Diabet Care*. 2008;**31**(1):105–7. doi: [10.2337/dc07-1154](https://doi.org/10.2337/dc07-1154). [PubMed: [17934145](https://pubmed.ncbi.nlm.nih.gov/17934145/)].
- Iseases. *Program on the Global Demography of Aging*. 2012. Available from: <https://EconPapers.repec.org/RePEc:gdms:paper:8712>.
- Roy T, Lloyd CE. Epidemiology of depression and diabetes: A systematic review. *J Affectiv Disorder*. 2012;**142**:S8–S21. doi: [10.1016/s0165-0327\(12\)70004-6](https://doi.org/10.1016/s0165-0327(12)70004-6).
- Moussavi S, Chatterji S, Verdes E, Tandon A, Patel V, Ustun B. Depression, chronic diseases, and decrements in health: Results from the World Health Surveys. *Lancet*. 2007;**370**(9590):851–8. doi: [10.1016/s0140-6736\(07\)61415-9](https://doi.org/10.1016/s0140-6736(07)61415-9).
- Corrigan JM, Greiner AC, Adams K. *1st Annual Crossing the Quality Chasm Summit: A focus on communities*. Washington, D.C: National Academies Press; 2004.
- Prince SA, Reed JL, Nerenberg KA, Kristjansson EA, Hiremath S, Adamo KB, et al. Intrapersonal, social and physical environmental determinants of moderate-to-vigorous physical activity in working-age women: A systematic review protocol. *Syst Rev*. 2014;**3**:132. doi: [10.1186/2046-4053-3-132](https://doi.org/10.1186/2046-4053-3-132). [PubMed: [25370736](https://pubmed.ncbi.nlm.nih.gov/25370736/)]. [PubMed Central: [PMC4224591](https://pubmed.ncbi.nlm.nih.gov/PMC4224591/)].
- Niknami M, Esmaeeli F, Nazer A, Ghoogazadeh M. Radiographic Evaluation of location and shape of mental foramen in selective Iranian Population. *J Dent Res*. 2011;**8**(2):102–8.
- Sniehotta FF, Scholz U, Schwarzer R. Bridging the intention-behaviour gap: Planning, self-efficacy, and action control in the adoption and maintenance of physical exercise. *Psychol Health*. 2005;**20**(2):143–60. doi: [10.1080/08870440512331317670](https://doi.org/10.1080/08870440512331317670).
- Widdershoven J, Kessing D, Schiffer A, Denollet J, Kupper N. How are depression and type D personality associated with outcomes in chronic heart failure patients? *Curr Heart Fail Rep*. 2013;**10**(3):244–53. doi: [10.1007/s11897-013-0139-7](https://doi.org/10.1007/s11897-013-0139-7). [PubMed: [23661199](https://pubmed.ncbi.nlm.nih.gov/23661199/)].
- Swenson SL, Rose M, Vittinghoff E, Stewart A, Schillinger D. The influence of depressive symptoms on clinician-patient communication among patients with type 2 diabetes. *Med Care*. 2008;**46**(3):257–65. doi: [10.1097/MLR.0b013e31816080e9](https://doi.org/10.1097/MLR.0b013e31816080e9). [PubMed: [18388840](https://pubmed.ncbi.nlm.nih.gov/18388840/)].

13. Khamseh ME, Baradaran HR, Rajabali H. Depression and diabetes in Iranian patients: A comparative study. *Int J Psychiatr Med*. 2007;**37**(1):81-6. doi: [10.2190/FP64-82V3-1741-842V](https://doi.org/10.2190/FP64-82V3-1741-842V). [PubMed: [17645200](https://pubmed.ncbi.nlm.nih.gov/17645200/)].
14. Rahimi A, Ahmadi F, Gholyaf M. The effects of Continuous Care Model on depression, anxiety, and stress in patients on hemodialysis. *Nephrol Nurs J*. 2008;**35**(1):39-43. [PubMed: [18372762](https://pubmed.ncbi.nlm.nih.gov/18372762/)].
15. Abbas Tavallaii S, Ebrahimnia M, Shamspour N, Assari S. Effect of depression on health care utilization in patients with end-stage renal disease treated with hemodialysis. *Eur J Intern Med*. 2009;**20**(4):411-4. doi: [10.1016/j.ejim.2009.03.007](https://doi.org/10.1016/j.ejim.2009.03.007). [PubMed: [19524185](https://pubmed.ncbi.nlm.nih.gov/19524185/)].
16. Sharif F, Shoul A, Janati M, Kojuri J, Zare N. The effect of cardiac rehabilitation on anxiety and depression in patients undergoing cardiac bypass graft surgery in Iran. *BMC Cardiovasc Disord*. 2012;**12**:40. doi: [10.1186/1471-2261-12-40](https://doi.org/10.1186/1471-2261-12-40). [PubMed: [22682391](https://pubmed.ncbi.nlm.nih.gov/22682391/)]. [PubMed Central: [PMC3420238](https://pubmed.ncbi.nlm.nih.gov/PMC3420238/)].
17. Lorig K, Stewart A, Ritter P, González V, Laurent D, Lynch J. *Outcome measures for health education and other health care interventions*. Thousand Oaks, USA: Sage; 1998.
18. Daniali SS, Eslami AA, Maracy MR, Shahabi J, Mostafavi-Darani F. The impact of educational intervention on self-care behaviors in overweight hypertensive women: A randomized control trial. *ARYA Atheroscler*. 2017;**13**(1):20-8. [PubMed: [28761451](https://pubmed.ncbi.nlm.nih.gov/28761451/)]. [PubMed Central: [PMC5515187](https://pubmed.ncbi.nlm.nih.gov/PMC5515187/)].
19. Kroenke K, Spitzer RL. The PHQ-9: A new depression diagnostic and severity measure. *Psychiatr Ann*. 2002;**32**(9):509-15. doi: [10.3928/0048-5713-20020901-06](https://doi.org/10.3928/0048-5713-20020901-06).
20. González VM, Stewart A, Ritter PL, Lorig K. Translation and validation of arthritis outcome measures into Spanish. *Arthrit Rheumatism*. 1995;**38**(10):1429-46. doi: [10.1002/art.1780381010](https://doi.org/10.1002/art.1780381010).
21. Wang Y, Lopez JM, Bolge SC, Zhu VJ, Stang PE. Depression among people with type 2 diabetes mellitus, US National Health and Nutrition Examination Survey (NHANES), 2005-2012. *BMC Psychiatry*. 2016;**16**:88. doi: [10.1186/s12888-016-0800-2](https://doi.org/10.1186/s12888-016-0800-2). [PubMed: [27044315](https://pubmed.ncbi.nlm.nih.gov/27044315/)]. [PubMed Central: [PMC4820858](https://pubmed.ncbi.nlm.nih.gov/PMC4820858/)].
22. Wittayanukorn S, Qian J, Hansen RA. Prevalence of depressive symptoms and predictors of treatment among U.S. adults from 2005 to 2010. *Gen Hosp Psychiatry*. 2014;**36**(3):330-6. doi: [10.1016/j.genhosppsych.2013.12.009](https://doi.org/10.1016/j.genhosppsych.2013.12.009). [PubMed: [24462337](https://pubmed.ncbi.nlm.nih.gov/24462337/)].
23. Gonzalez JS, Safren SA, Cagliero E, Wexler DJ, Delahanty L, Wittenberg E, et al. Depression, self-care, and medication adherence in type 2 diabetes: relationships across the full range of symptom severity. *Diabetes Care*. 2007;**30**(9):2222-7. doi: [10.2337/dc07-0158](https://doi.org/10.2337/dc07-0158). [PubMed: [17536067](https://pubmed.ncbi.nlm.nih.gov/17536067/)]. [PubMed Central: [PMC4440862](https://pubmed.ncbi.nlm.nih.gov/PMC4440862/)].
24. Egede LE, Ellis C, Grubaugh AL. The effect of depression on self-care behaviors and quality of care in a national sample of adults with diabetes. *Gen Hosp Psychiatry*. 2009;**31**(5):422-7. doi: [10.1016/j.genhosppsych.2009.06.007](https://doi.org/10.1016/j.genhosppsych.2009.06.007). [PubMed: [19703635](https://pubmed.ncbi.nlm.nih.gov/19703635/)].
25. Schoenthaler A, Ogedegbe G, Allegrante JP. Self-efficacy mediates the relationship between depressive symptoms and medication adherence among hypertensive African Americans. *Health Educ Behav*. 2009;**36**(1):127-37. doi: [10.1177/1090198107309459](https://doi.org/10.1177/1090198107309459). [PubMed: [18077654](https://pubmed.ncbi.nlm.nih.gov/18077654/)].
26. Egede LE, Ellis C. Diabetes and depression: Global perspectives. *Diabetes Res Clin Pract*. 2010;**87**(3):302-12. doi: [10.1016/j.diabres.2010.01.024](https://doi.org/10.1016/j.diabres.2010.01.024). [PubMed: [20181405](https://pubmed.ncbi.nlm.nih.gov/20181405/)].
27. Jacka FN, Pasco JA, Mykletun A, Williams LJ, Hodge AM, O'Reilly SL, et al. Association of Western and traditional diets with depression and anxiety in women. *Am J Psychiatry*. 2010;**167**(3):305-11. doi: [10.1176/appi.ajp.2009.09060881](https://doi.org/10.1176/appi.ajp.2009.09060881). [PubMed: [20048020](https://pubmed.ncbi.nlm.nih.gov/20048020/)].
28. Bair MJ, Robinson RL, Katon W, Kroenke K. Depression and pain comorbidity: A literature review. *Arch Intern Med*. 2003;**163**(20):2433-45. doi: [10.1001/archinte.163.20.2433](https://doi.org/10.1001/archinte.163.20.2433). [PubMed: [14609780](https://pubmed.ncbi.nlm.nih.gov/14609780/)].
29. Kav S, Yilmaz AA, Bulut Y, Dogan N. Self-efficacy, depression and self-care activities of people with type 2 diabetes in Turkey. *Collegian*. 2017;**24**(1):27-35. [PubMed: [29218959](https://pubmed.ncbi.nlm.nih.gov/29218959/)].
30. Kaur G, Tee GH, Ariaratnam S, Krishnapillai AS, China K. Depression, anxiety and stress symptoms among diabetics in Malaysia: A cross sectional study in an urban primary care setting. *BMC Fam Pract*. 2013;**14**:69. doi: [10.1186/1471-2296-14-69](https://doi.org/10.1186/1471-2296-14-69). [PubMed: [23710584](https://pubmed.ncbi.nlm.nih.gov/23710584/)]. [PubMed Central: [PMC3669622](https://pubmed.ncbi.nlm.nih.gov/PMC3669622/)].
31. Oliveira A, Sá M, Freitas A. Learning to Live with a Chronic Illness, without Thinking about Death. *Academ J Interdisciplin Stud*. 2013;**2**(9):326. doi: [10.5901/ajis.2013.v2n9p326](https://doi.org/10.5901/ajis.2013.v2n9p326).
32. Khuwaja AK, Lalani S, Dhanani R, Azam IS, Rafique G, White F. Anxiety and depression among outpatients with type 2 diabetes: A multi-centre study of prevalence and associated factors. *Diabetol Metab Syndr*. 2010;**2**:72. doi: [10.1186/1758-5996-2-72](https://doi.org/10.1186/1758-5996-2-72). [PubMed: [21171976](https://pubmed.ncbi.nlm.nih.gov/21171976/)]. [PubMed Central: [PMC3022608](https://pubmed.ncbi.nlm.nih.gov/PMC3022608/)].
33. Van de Velde S, Bracke P, Levecque K. Gender differences in depression in 23 European countries. Cross-national variation in the gender gap in depression. *Soc Sci Med*. 2010;**71**(2):305-13. doi: [10.1016/j.socscimed.2010.03.035](https://doi.org/10.1016/j.socscimed.2010.03.035). [PubMed: [20483518](https://pubmed.ncbi.nlm.nih.gov/20483518/)].