

Phytochemicals in Cancer Prevention: A Review of the Evidence

Neda Mollakhalili Meybodi,^{1,2} Amir Mohammad Mortazavian,^{3,*} Ayad Bahadori Monfared,^{4,*} Sara

Sohrabvandi,⁵ and Fateme Aghaei Meybodi⁶

¹Student Research Office, Department of Food Science and Technology, Faculty of Nutrition Sciences and Food Technology/National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

²Research Center for Molecular Identification of Food Hazards, Shahid Sadoughi University of Medical Sciences, Yazd, IR Iran

³Department of Food Science and Technology, Faculty of Nutrition Sciences, Food Science and Technology/National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

⁴Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

⁵Department of Food Technology Research, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

⁶Department of Internal Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, IR Iran

*Corresponding authors: Amir Mohammad Mortazavian, PhD, Department of Food Science and Technology, Faculty of Nutrition Sciences, Food Science and Technology/National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, P.O. Box: 19395-4741, Tehran, IR Iran. E-mail: mortazavn@sbmu.ac.ir; mortazavn@yahoo.com; Ayad Bahadori Monfared, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, P.O. Box: 19395-4741, Tehran, IR Iran. E-mail: a.bahadorimonfared@sbmu.ac.ir

Received 2016 May 25; Revised 2016 August 19; Accepted 2017 January 07.

Abstract

Context: Cancer generally is considered as a neoplastic disease with particular causative and etiologic factors as well as protective elements. Although it has remained difficult to treat, it is preventable. Recently, the interest in dietary phytochemicals intake has considerably increased for potential cancer chemoprevention.

Evidence Acquisition: This report reviews the role of phytochemical consumption in cancer prevention based on publications from PubMed, Science direct, Google Scholar, and Scopus from the year 1996 onward using cancer, chemoprevention, phytochemical keywords.

Results: Regular intake of phytochemicals has been demonstrated to prevent cancer during its different stages including initiation, promotion and progression. Considering the animal models, the second step is the main stage for cancer chemoprevention.

Conclusions: The phytochemicals involved in chemoprevention can be categorized in different groups naming phenolics, carotenoid, alkaloids, organosulfur compounds and nitrogen-containing compounds. They are able to stop, postpone and reverse carcinogenesis by different mechanisms.

Keywords: Cancer, Chemoprevention, Phytochemical

1. Introduction

Cancer as a genetic disorder is the main cause of death in economically developed countries and the second one in emerging countries (1). The reports indicate that about 13% of total deaths (7.6 million) are induced by cancer and its global burden is increased largely regarding both the aging and growth of the world population besides the growing of cancer inducing behaviors, especially smoking (2, 3). The majority of these disorders are generated by lung, stomach, liver, colon and breast cancer (4). It has been estimated that about 50% - 60% of cancer enduring in the United States exploit compounds deriving from different parts of plants or nutrients (as a complementary agent and/or substitute medicine), solely or alongside with traditional therapeutic treatment like chemotherapy and/or radiation therapy (5-7).

Natural herbs have been used for thousands of years for prevention and/or treatment of diverse diseases (8). The presence of bioactive components in plants makes

them appropriate choices to be used especially by gourmet food consumers (9-11). Evidences confirmed the anticancer activities of natural plants derived bioactive components (12-16). The bulk of research has been devoted to dietary phytochemicals which resulted in an increase in comprehension of these compounds as a chemical and biological functional agent which has a constructive effect on human health. Regarding to development of studies which are done in vitro (advent of different cellular, molecular, and genomic trial systems) and in vivo (transgenic and knockout animal models), the mechanisms by which dietary phytochemicals are involved has been greatly understood.

Upon entering cells, the phytochemicals are able to hunt free radicals immediately (10, 17) and create the signals in response to chemical or electrophilic stress that activate proteins associated to diverse cellular signaling pathways (18, 19). The mentioned ability involves the activation of the nuclear factor erythroid-2 (NF-E2)-related factor 2 (Nrf2)-Kelch-like ECH associated protein 1 (Keap1)

(Nrf2-Keap1 complex) (20, 21). This complex activation consequently influences the cellular defense mechanisms, counting phase II detoxifying enzymes, phase III transporters, anti-oxidative stress proteins, and other stress-defense molecules induction which defend the normal cells against reactive oxygen species (ROS), reactive nitrogen species (RNS) and/or reactive metabolites of carcinogenic species (22, 23). They are also able to cause death of apoptotic cell in pre-neoplastic or neoplastic cells via different growth suppression methods comprising the organization of cytochrome c (Cyt c)/caspases, cell cycle catch, the embarrassment of the nuclear factor- κ B (NF- κ B), Janus kinase (JAK)-signal transducer, and activator of transcription (STAT) signaling pathways, resulting in the tumor progression reticence (24, 25). It is worthy to consider that advanced/metastatic cancers which have highly genetic mutations, loss of heterozygosity, and/or epigenetic changes are greatly resistant even toward radiation or chemotherapeutic drugs and so they will not be responded to dietary phytochemicals alone (26, 27). These defensive mechanisms that prevent the induction of carcinogenesis are defined as chemoprevention perception that was originally established by Wattenberg (1966) (28, 29).

Chemoprevention passes on using agents able to prevent, reverse or postpone tumorigenesis (15, 30). It has been reported that several phytochemicals originated of edible plants are able to impede with a specific stage of the carcinogenic process. Different studies revealed that phytochemicals are able to be a chemopreventative agent toward the human cancer by inflection of the cancer cell cycle, proliferation inhibition, and initiation of apoptosis (31). Considering both the cancer epidemiology and trial finding it has been revealed that dietary phytochemicals can be used rightly in chemoprevention and its daily consumption is a hopeful new attitude to avoid carcinogenesis. In this review, an attempt has been devoted to introduce dietary phytochemicals, their healthful impacts to prevent cancer along with their mechanisms of action.

2. Evidence Acquisition

Review was conducted using keywords such as cancer, chemoprevention, phytochemical thorough PubMed, Science direct, Google Scholar, and Scopu sites. High quality articles and books were reviewed. Articles and books providing constructive information to the topic were further comprised in the current study.

3. Results

3.1. Phytochemicals

Plant derived phytochemicals are defined as bioactive non-nutrient compounds which have been connected to reduction of major chronic diseases risk (32-34). The Greek word 'phyto' in phytochemicals means plant (35, 36). In other words, phytochemicals are plant chemicals. It is predicted that more than 5000 particular phytochemicals have been recognized in grains, fruits and vegetables but a large percentage are still unknown and must be identified before understanding their health benefits in whole foods (10, 37). An ideal chemopreventive factor must be considered by different criteria as mentioned in Figure 1.




Figure 1. The Characteristics of an Ideal Chemopreventive Agent

Phytochemicals are able to impede initiation or repeal the promotion step of multistep carcinogenesis (15, 38). They can also stop or postpone the development of pre-cancerous cells into the malignant ones (39, 40). Nevertheless, evidences indicated that the health benefits of phytochemicals in fruits and vegetables would be even greater than it is currently implicit, regarding the role of oxidative stress induced free radicals in a wide range of chronic diseases.

Different biologically active phytochemicals have been identified to have the capability to control the carcinogenesis at different stages which are presented below and summarized in Figure 2.




Figure 2. Classification of Dietary Phytochemicals

3.1.1. Phenolics

The secondary metabolites phenolics are extensively found in fruits (41). These compounds have one or more aromatic rings with one or more hydroxyl groups (42). The main phenolic compounds which are found in food could be categorized into three groups: simple phenols and phenolic acids, hydroxycinnamic acid derivatives and flavonoids. They play an essential role in the growth and reproduction of the plants besides acting as defense mechanisms against pathogens, parasites, and predators (43, 44). The upward attention in these compounds is mainly attributed to their antioxidant capacity and their ability to prevent some diseases. It is notable that the phytochemicals health benefits will be achieved by their regular intake and their bioavailability (45).

Simple phenols and phenolic acids are precursor to the synthesis of other complex compounds such as flavonoids and tannins (46, 47). They act in the natural defense mechanism of plants to prevent them from infectious diseases and reduce the growth of pathogenic bacteria, viruses, and fungi. They include monophenols, 3-ethylphenol, 3, 4-dimethylphenol and diphenols which are possibly the most prevalent simple phenol (47).

Hydroxycinnamic acids and their esterified derivatives in fruits and vegetables are almost completely originated from p-coumaric acid (PCA), caffeic acid (CA), and ferulic acid (FA) (48). They are frequently present in conjugated forms, usually as esters rather than glycosides (49). Hydroxycinnamic acids derivatives like chlorogenic

acid, coumaric acid, caffeic acid and sinapic acid are the foremost of phytochemicals in the plant food powders (50). Consequently, the addition of these food plant powders high in bioactive phytochemicals will beneficially create food reformulations able to improve foods regarding both the quality and health endorsement properties. This approach will directly improve the plant-based foods consumption which is established as being beneficial for health.

Flavonoids as a major class of phenolic compounds exhibited highly antioxidant activity (51). These compounds have been connected to reducing the risk of main chronic diseases and have been recognized largely in fruits, vegetables, and other plant foods (52). More than 4000 diverse flavonoids have been known. They frequently have a general structure containing two aromatic rings (A and B rings) connected via 3 carbons which are typically in an oxygenated heterocycle ring or C rings. Flavonols, flavones, flavanols (catechins), flavanones, anthocyanidins, and isoflavonoids are different types of flavonoids considering the variation of the general structure of the heterocyclic C ring (53).

3.1.2. Organosulfur Compounds

Organosulfur compounds are organic compounds which can be recognized according to their sulfur containing functional groups (54). The regular intake of organosulfur compounds imparts bioactive properties especially about cardiovascular health (37). Several organosulfur compounds were assessed by Wattenberg et al considering their ability to prevent carcinogenesis stimulated by N-nitrosodiethylamine, and the strongest was diallyl disulfide (55). Considering the ability of natural products derived from cruciferous plants and members of *Allium* genus to prevent cancers, these vegetables have been investigated in several studies. Watercress, Chinese cabbage and broccoli are some examples of vegetable rich in organosulfur compounds (OSCs).

3.1.3. Carotenoids

Carotenoids as the most extensive natural pigment have gained considerable attention regarding their provitamin and antioxidant properties (56, 57). More than 600 different carotenoids have been recognized in nature. They may be of plants, microorganisms, and/or animals origin. They have a skeleton of isoprene units containing 40-carbon (58). Their structure may be cyclized at one or both ends, having diverse hydrogenation degree, holding oxygen-containing functional groups. Carotenoid compounds are mainly present in all-trans form in nature. The most typical properties of carotenoids are their long sequences of conjugated double bonds formed at the cen-

ter of the molecule. This characteristic is impressive in their shape, chemical reactivity, and light-absorbing properties. Carotenoids are able to react with free radicals and create radicals themselves (59). The existence of sufficient carotenoids can inhibit lipid oxidation and related oxidative stress.

3.1.4. Alkaloids

Alkaloids are a group of ring structure nitrogen containing organic compounds with a wide range of anti-cancer activity (60). These compounds take part in cancer inhibition via prevention of enzyme topoisomerase activity which is involved in DNA imitation, inducing apoptosis and expression of p53 gene (61, 62). However, alkaloids have long been existed before humans; some of them are structurally similar to neurotransmitters present in the central nervous system of humans. Considering the medicinal importance of alkaloids and research explaining their role in treating wild propagation of cells, they would be used as an effective chemopreventive agent in the era of modern drug innovation (63, 64). Amaryllidaceae alkaloids, betalain alkaloids, diterpenoid alkaloids, indole alkaloids, isoquinoline alkaloids, lycopodium alkaloids, monoterpene and sesquiterpene alkaloids, peptide alkaloids, pyrrolidine and piperidine alkaloids, pyrrolizidine alkaloids, quinoline alkaloids, quinolizidine alkaloids, steroidal alkaloids, tropane alkaloids, and miscellaneous alkaloids are the main categories of Alkaloids.

3.2. Cancer Chemoprevention by Phytochemicals

Cancer, a multistage, multi-mechanism carcinogenesis process, comprises mutagenic, cell death and epigenetic mechanisms, through three separate but closely linked stages: initiation, promotion, and progression (65, 66). The first step, initiation, is mainly concluded from a single using of a subcarcinogenic dose of a carcinogen (67). In another words, exposing to carcinogen agents creates permanent injuries to genetic material which are almost irreversible. Promotion, the second step, resulted via repeatedly applying of an irritating agent (68). It involves cellular explosion and selective clonal growth which are reversible, during its early stages, but becomes irreversible with time. Cells in humans and other organisms are frequently interpretate to a variety of oxidizing agents which are necessary for life in some cases (69, 70). These factors may be present in air, food, and water, or may be created during metabolic activity within cells (43). The main factor that must be considered is keeping equilibrium between oxidants and antioxidants to prolong the optimal physiological conditions. An imbalance created by producing high amounts of oxidants, will, lead to oxidative stress,

especially in chronic bacterial, viral, and parasitic infections (10). Oxidative stress can hurt large biomolecules like lipids, proteins, and DNA, resulting in an increased risk for cancer and cardio-vascular diseases (CVD). Since the initiation phase reduction to a zero level is impossible, the main interference would be achieved at the promotion level to eradicate premalignant cells before being malignant (50). The conversion of the normal cells to malignant ones took place during several years. So, their delaying or prevention of this transformation is a viable and possible objective for the future (70). The results of many laboratory animal studies evidently denote that different cancers can be prohibited using certain chemicals. In order to avoid and/or slow the oxidative stress created by free radicals, adequate amounts of antioxidants are necessary to be used. Cancer chemoprevention is the main cancer preventive approach that exploits naturally dietary phytochemicals or remedial drugs with fairly low toxicity. The anti-cancer properties of phytochemicals are presented in Table 1. Different mechanisms involved in chemoprevention of different phytochemicals are presented in Table 2.

Table 1. The Anticancer Properties of Phytochemicals

Phytochemicals	Anti-Cancer Properties
Phenolic compound	Reduced incidence of neoplasia induced by chemical carcinogens
	Preventing nitrosation of susceptible secondary amines and amides to form highly potent carcinogenic nitrosamines and nitrosamides in our foods
	Potent chemical nucleophiles
	Inhibitors of promotion processes
	Inhibitors of kinases by reducing hyperproliferation of Epithelial cells
Organosulfur compounds	Induction of carcinogen detoxification
	Inhibition of tumor cell proliferation
	Antimicrobial effect
	Free radical scavenging
	Inhibition of DNA adduct formation
	Induction of cell cycle arrest
Alkaloids	Induction of apoptosis
	Modification of carcinogen metabolism
	Modification of tumor metabolism
Carotenoids	Inhibition of tumor cell growth
	Inducers of differentiation
Nitrogen containing compounds	Inhibit the metabolic activation and carcinogenicity

As is presented in Table 2 the success of chemopreventive agents is dependent on their ability to neutralize the precise upstream signals which create different forms of cellular stress, genotoxic damage and redox imbalances.

Different phytochemicals in fruits and vegetables have been detached and recognized which their ability to im-

Table 2. The Mechanistic Insight Into Chemoprevention of Different Phytochemicals

Mechanism of Cancer Prevention	Reference
Antioxidant activity	
Scavenge free radicals and reduce oxidative stress	
Inhibition of	
Cell proliferation	
Cell differentiation	
Oncogene expression	
Signal transduction pathways	
Induction of	
Tumor suppress gene expression	
cell-cycle arrest	
Enzyme induction and enhancing detoxification	
Phase II enzyme	
Glutathione peroxidase	
Catalase	(9)
Superoxide dismutase	
Enzyme inhibition	
Phase I enzyme (block activation of carcinogens)	
Cyclooxygenase-2	
Inducible nitric oxide synthase	
Xanthine oxide	
Enhancement of immune functions and surveillance	
Antiangiogenesis	
Inhibition of cell adhesion and invasion	
Inhibition of nitrosation and nitration	
Prevention of DNA binding	
Regulation of steroid hormone metabolism	
Regulation of estrogen metabolism	
Antibacterial and antiviral effects	

pede different stages of the carcinogenic process in numerous animal models have been demonstrated (52, 58, 61). Chemicals which are able to avoid the development of carcinogens from precursor substances or to inhibit carcinogens from attaining or responding to critical target DNA sites in the tissues are named 'blocking agents'. Chemicals which keep on suppressing the illustration of neoplasia in cells interpretate to doses of a carcinogenic agent are called 'suppressing agents'. Administration of vegetables and/or fruits or their components in the diet to animals is able to reduce chemically-generated tumor occurrence (53). As previously mentioned, carcinogenic species, like environmental pollutants, dietary mutagens and radiation, will possibly result in the creation of reactive oxygen species (ROS) and/or reactive nitrogen species (RNS), which react with cellular molecules like proteins, lipids, and DNA to provoke carcinogenesis. Dietary phytochemicals intake not only scavenge ROS/RNS directly but also

eliminate carcinogenic reactive intermediates indirectly by the transcription factor Nrf2 [nuclear factor erythroid 2 p45 (NF-E2)-related factor 2] antioxidant and detoxification system. When Nrf2 is released from Kelch-like ECH associated protein 1 (Keap1) and translocates to the nucleus, Nrf2 binds to antioxidant responsive elements (AREs) in the promoter/enhancer region of phase II detoxification and antioxidant enzyme genes with the Maf protein. Recent research has also shown that the reactivation of Nrf2 might be regulated by dietary phytochemicals through epigenetic modifications such as DNA methylation and histone modification.

Block et al. investigated more than 200 epidemiological cases that studied the daily intake of fruits and vegetables effects on different cancers naming lung, colon, breast, cervix, esophagus, oral cavity, stomach, bladder, pancreas, and ovary (36). In 128 of 156 dietary studies, utilization of fruits and vegetables create a significant protective effect. Results indicated that the cancer risk decreased due to consumption fruits and vegetables. Significant protection was observed in 24 of 25 studies for lung cancer (35). Fruits consumption was significantly able to create a protective effect in the case of esophagus, oral cavity, and larynx cancers. Their intake was also protective for cancer of the pancreas and stomach in 26 of 30 studies and for colorectal and bladder cancer in 23 of 38 studies. Studies to date which demonstrated the ability of common phytochemicals regarding the cancer which they are preventive are presented in Table 3.

4. Conclusions

The process of carcinogenesis is complex and heterogeneous regarding to several combinations of genetic and epigenetic events which occur in an individual cell to create a neoplastic deformation. Considering different stages of cancer, initiation, promotion and progression, the second step is the main one to be considered for cancer chemoprevention. It has been predicted that more than two-thirds of human cancers could be prevented via proper lifestyle adjustment. Phytochemicals are mainly present in fruits and vegetables and their regular intake would be effective to cancer prevention.

Acknowledgments

We are grateful to Shahid Beheshti University of medical science for support of this study.

Table 3. The ability of Common Phytochemicals Regarding Their Cancer Preventive Effects

Phytochemicals	Cancer Target	Results	Reference
Cranberry extracts	Breast cancer MCF-7 cells	Cranberry phytochemical extracts possess the ability to suppress the proliferation of human breast cancer MCF-7 cells	(6)
Blueberry extract	MDA-MB-231 Breast Cancer Cells	Decreased phosphatidylinositol 3-kinase (PI3K)/AKT and NF- κ B activation in MDA-MB-breast cancer cells	(18)
Carotenoids, retinol, and tocopherols	Breast cancer	The results indicated an inverse association of carotenoids and breast cancer among postmenopausal women	(39)
Selenium and Vitamin E	Prostate cancer	Dietary supplementation with vitamin E significantly increased the risk of prostate cancer among healthy men	(25)
Glucosinolates (secondary metabolites produced by crucifers)	Prostate cancer	Inhibit prostate cancer by both blocking initiation and suppressing prostate cancer progression in vitro and in vivo	(42)
Quercetin	Cancer	Quercetin can be efficient at treating cancer by inducing cell death or cell cycle arrest preferentially in cancer cells versus their normal counterparts through a process involving the down-regulation of selective oncogenes (such as Mcl-1, Ras, MEK, PI3K), or the up-regulation of tumor suppressor genes which, in turn enhance selective pathways leading to the elimination of cancer cells	(38)
Resveratrol (polyphenol)	Colorectal cancer, pancreatic cancer, stem cell breast cancer	Resveratrol induces cell apoptosis like as many other polyphenols	(47)
Tocopherols, tocotrienols and γ -oryzanol (rice bran extracts)	Colorectal cancer	Rice bran extracts differentially inhibit colorectal cancer (CRC) cell growth. Rice bran chemoprevention is due to the complex phytochemical mixture	(52)

Footnotes

Conflict of Interest: The authors have no conflict of interest in this study.

Authors' Contribution: Neda Mollakhalili Meybodi collected the data designed this article. Neda Mollakhalili Meybodi and Amir Mohammad Mortazavian participate in preparing of the manuscript. All authors read and approved the final manuscript.

Financial Disclosure: None declared.

References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin.* 2011;**61**(2):69–90. doi: [10.3322/caac.20107](https://doi.org/10.3322/caac.20107). [PubMed: [21296855](https://pubmed.ncbi.nlm.nih.gov/21296855/)].
- Zhang J. Can We Discover “Really Safe and Effective” Anticancer Drugs?. *Adv Pharmacoevidiol Drug Safety.* 2012;**01**(05) doi: [10.4172/2167-1052.1000e113](https://doi.org/10.4172/2167-1052.1000e113).
- Schottenfeld D, Beebe-Dimmer JL, Buffler PA, Omenn GS. Current perspective on the global and United States cancer burden attributable to lifestyle and environmental risk factors. *Annu Rev Public Health.* 2013;**34**:97–117. doi: [10.1146/annurev-publhealth-031912-114350](https://doi.org/10.1146/annurev-publhealth-031912-114350). [PubMed: [23514316](https://pubmed.ncbi.nlm.nih.gov/23514316/)].
- Chen W, Zheng R, Zhang S, Zhao P, Zeng H, Zou X, et al. Annual report on status of cancer in China, 2010. *Chin J Cancer Res.* 2014;**26**(1):48–58. doi: [10.3978/j.issn.1000-9604.2014.01.08](https://doi.org/10.3978/j.issn.1000-9604.2014.01.08). [PubMed: [24653626](https://pubmed.ncbi.nlm.nih.gov/24653626/)].
- A. Ajith T, K. Janardhanan K. Indian medicinal mushrooms as a source of antioxidant and antitumor agents. *J Clin Biochem Nutr.* 2007;**40**(3):157–62. doi: [10.3164/jcfn.40.157](https://doi.org/10.3164/jcfn.40.157). [PubMed: [18398492](https://pubmed.ncbi.nlm.nih.gov/18398492/)].
- Sun J, Hai Liu R. Cranberry phytochemical extracts induce cell cycle arrest and apoptosis in human MCF-7 breast cancer cells. *Cancer Lett.* 2006;**241**(1):124–34. doi: [10.1016/j.canlet.2005.10.027](https://doi.org/10.1016/j.canlet.2005.10.027). [PubMed: [16377076](https://pubmed.ncbi.nlm.nih.gov/16377076/)].
- Lawenda BD, Kelly KM, Ladas EJ, Sagar SM, Vickers A, Blumberg JB. Should supplemental antioxidant administration be avoided during chemotherapy and radiation therapy?. *J Natl Cancer Inst.* 2008;**100**(11):773–83. doi: [10.1093/jnci/djn148](https://doi.org/10.1093/jnci/djn148). [PubMed: [18505970](https://pubmed.ncbi.nlm.nih.gov/18505970/)].
- Lai PK, Roy J. Antimicrobial and chemopreventive properties of herbs and spices. *Curr Med Chem.* 2004;**11**(11):1451–60. [PubMed: [15180577](https://pubmed.ncbi.nlm.nih.gov/15180577/)].
- Temple NJ, Wilson T, Jacobs Jr DR. Nutritional health: strategies for disease prevention. Springer Science & Business Media; 2012.
- Liu RH. Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals. *Am J Clin Nutr.* 2003;**78**(3 Suppl):517S–20S. [PubMed: [12936943](https://pubmed.ncbi.nlm.nih.gov/12936943/)].
- Wang L, Weller CL. Recent advances in extraction of nutraceuticals from plants. *Trends Food Sci Technol.* 2006;**17**(6):300–12. doi: [10.1016/j.tifs.2005.12.004](https://doi.org/10.1016/j.tifs.2005.12.004).
- Pezzuto JM. Plant-derived anticancer agents. *Biochem Pharmacol.* 1997;**53**(2):121–33. [PubMed: [9037244](https://pubmed.ncbi.nlm.nih.gov/9037244/)].
- Shukla Y, Singh M. Cancer preventive properties of ginger: a brief review. *Food Chem Toxicol.* 2007;**45**(5):683–90. doi: [10.1016/j.fct.2006.11.002](https://doi.org/10.1016/j.fct.2006.11.002). [PubMed: [17175086](https://pubmed.ncbi.nlm.nih.gov/17175086/)].
- Aggarwal BB, Kunnumakkara AB, Harikumar KB, Tharakan ST, Sung B, Anand P. Potential of spice-derived phytochemicals for cancer prevention. *Planta Med.* 2008;**74**(13):1560–9. doi: [10.1055/s-2008-1074578](https://doi.org/10.1055/s-2008-1074578). [PubMed: [18612945](https://pubmed.ncbi.nlm.nih.gov/18612945/)].
- Surh YJ. Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer.* 2003;**3**(10):768–80. doi: [10.1038/nrc1189](https://doi.org/10.1038/nrc1189). [PubMed: [14570043](https://pubmed.ncbi.nlm.nih.gov/14570043/)].
- Mans DR, da Rocha AB, Schwartzmann G. Anti-cancer drug discovery and development in Brazil: targeted plant collection as a rational strategy to acquire candidate anti-cancer compounds. *Oncologist.* 2000;**5**(3):185–98. [PubMed: [10884497](https://pubmed.ncbi.nlm.nih.gov/10884497/)].
- Johns T. Phytochemicals as Evolutionary Mediators of Human Nutritional Physiology. *Pharm Biol.* 1996;**34**(5):327–34. doi: [10.1076/phbi.34.5.327.13254](https://doi.org/10.1076/phbi.34.5.327.13254).
- Adams LS, Phung S, Yee N, Seeram NP, Li L, Chen S. Blueberry phytochemicals inhibit growth and metastatic potential of MDA-MB-231 breast cancer cells through modulation of the phosphatidylinositol 3-kinase pathway. *Cancer Res.* 2010;**70**(9):3594–605. doi: [10.1158/0008-5472.CAN-09-3565](https://doi.org/10.1158/0008-5472.CAN-09-3565). [PubMed: [20388778](https://pubmed.ncbi.nlm.nih.gov/20388778/)].
- Finley JW, Kong AN, Hintze KJ, Jeffery EH, Ji LL, Lei XG. Antioxidants in foods: state of the science important to the food industry. *J Agric Food Chem.* 2011;**59**(13):6837–46. doi: [10.1021/jf2013875](https://doi.org/10.1021/jf2013875). [PubMed: [21627162](https://pubmed.ncbi.nlm.nih.gov/21627162/)].
- Kim J, Cha YN, Surh YJ. A protective role of nuclear factor-erythroid 2-related factor-2 (Nrf2) in inflammatory disorders. *Mutat Res.* 2010;**690**(1-2):12–23. doi: [10.1016/j.mrfmmm.2009.09.007](https://doi.org/10.1016/j.mrfmmm.2009.09.007). [PubMed: [19799917](https://pubmed.ncbi.nlm.nih.gov/19799917/)].
- Surh YJ, Kundu JK, Na HK. Nrf2 as a master redox switch in turning on the cellular signaling involved in the induction of cytoprotective genes by some chemopreventive phytochemicals. *Planta Med.* 2008;**74**(13):1526–39. doi: [10.1055/s-0028-1088302](https://doi.org/10.1055/s-0028-1088302). [PubMed: [18937164](https://pubmed.ncbi.nlm.nih.gov/18937164/)].

22. Lee JH, Khor TO, Shu L, Su ZY, Fuentes F, Kong AN. Dietary phytochemicals and cancer prevention: Nrf2 signaling, epigenetics, and cell death mechanisms in blocking cancer initiation and progression. *Pharmacol Ther*. 2013;**137**(2):153-71. doi: [10.1016/j.pharmthera.2012.09.008](https://doi.org/10.1016/j.pharmthera.2012.09.008). [PubMed: [23041058](https://pubmed.ncbi.nlm.nih.gov/23041058/)].
23. Kumar H, Kim IS, More SV, Kim BW, Choi DK. Natural product-derived pharmacological modulators of Nrf2/ARE pathway for chronic diseases. *Nat Prod Rep*. 2014;**31**(1):109-39. doi: [10.1039/c3np70065h](https://doi.org/10.1039/c3np70065h). [PubMed: [24292194](https://pubmed.ncbi.nlm.nih.gov/24292194/)].
24. Seeram NP, Adams LS, Zhang Y, Lee R, Sand D, Scheuller HS, et al. Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells in vitro. *J Agric Food Chem*. 2006;**54**(25):9329-39. doi: [10.1021/jf061750g](https://doi.org/10.1021/jf061750g). [PubMed: [17147415](https://pubmed.ncbi.nlm.nih.gov/17147415/)].
25. Klein EA, Thompson IJ, Tangen CM, Crowley JJ, Lucia MS, Goodman PJ, et al. Vitamin E and the risk of prostate cancer: the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA*. 2011;**306**(14):1549-56. doi: [10.1001/jama.2011.1437](https://doi.org/10.1001/jama.2011.1437). [PubMed: [21990298](https://pubmed.ncbi.nlm.nih.gov/21990298/)].
26. Kaur M, Agarwal C, Agarwal R. Anticancer and cancer chemopreventive potential of grape seed extract and other grape-based products. *J Nutr*. 2009;**139**(9):1806S-12S. doi: [10.3945/jn.109.106864](https://doi.org/10.3945/jn.109.106864). [PubMed: [19640973](https://pubmed.ncbi.nlm.nih.gov/19640973/)].
27. Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med*. 2001;**344**(11):783-92. doi: [10.1056/NEJM200103153441101](https://doi.org/10.1056/NEJM200103153441101). [PubMed: [11248153](https://pubmed.ncbi.nlm.nih.gov/11248153/)].
28. Pereira MA, Grubbs CJ, Barnes LH, Li H, Olson GR, Eto I, et al. Effects of the phytochemicals, curcumin and quercetin, upon azoxymethane-induced colon cancer and 7,12-dimethylbenz[a]anthracene-induced mammary cancer in rats. *Carcinogenesis*. 1996;**17**(6):1305-11. [PubMed: [8681447](https://pubmed.ncbi.nlm.nih.gov/8681447/)].
29. Shu L, Cheung KL, Khor TO, Chen C, Kong AN. Phytochemicals: cancer chemoprevention and suppression of tumor onset and metastasis. *Cancer Metastasis Rev*. 2010;**29**(3):483-502. doi: [10.1007/s10555-010-9239-y](https://doi.org/10.1007/s10555-010-9239-y). [PubMed: [20798979](https://pubmed.ncbi.nlm.nih.gov/20798979/)].
30. Surh YJ, Na HK, Lee SS. Transcription factors and mitogen-activated protein kinases as molecular targets for chemoprevention with anti-inflammatory phytochemicals. *Biofactors*. 2004;**21**(1-4):103-8. [PubMed: [15630178](https://pubmed.ncbi.nlm.nih.gov/15630178/)].
31. Gonzalez de Mejia E, Song YS, Ramirez-Mares MV, Kobayashi H. Effect of yerba mate (*Ilex paraguariensis*) tea on topoisomerase inhibition and oral carcinoma cell proliferation. *J Agric Food Chem*. 2005;**53**(6):1966-73. doi: [10.1021/jf048158g](https://doi.org/10.1021/jf048158g). [PubMed: [15769122](https://pubmed.ncbi.nlm.nih.gov/15769122/)].
32. Wiseman H. The bioavailability of non-nutrient plant factors: dietary flavonoids and phyto-oestrogens. *Proc Nutr Soc*. 1999;**58**(1):139-46. [PubMed: [10343351](https://pubmed.ncbi.nlm.nih.gov/10343351/)].
33. Sreelatha S, Padma PR. Antioxidant activity and total phenolic content of Moringa oleifera leaves in two stages of maturity. *Plant Foods Hum Nutr*. 2009;**64**(4):303-11. doi: [10.1007/s11130-009-0141-0](https://doi.org/10.1007/s11130-009-0141-0). [PubMed: [19904611](https://pubmed.ncbi.nlm.nih.gov/19904611/)].
34. Doughari JH, Human IS, Benadé AJ, Nkaidemi PA. Phytochemicals as chemotherapeutic agents and antioxidants: Possible solution to the control of antibiotic resistant verocytotoxin producing bacteria. *Planta medica*. 2009;**3**(11):839-48.
35. Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr*. 2004;**134**(12 Suppl):3479S-85S. [PubMed: [15570057](https://pubmed.ncbi.nlm.nih.gov/15570057/)].
36. Block G, Norkus E, Hudes M, Mandel S, Helzlsouer K. Which plasma antioxidants are most related to fruit and vegetable consumption?. *Am J Epidemiol*. 2001;**154**(12):1113-8. [PubMed: [11744516](https://pubmed.ncbi.nlm.nih.gov/11744516/)].
37. Liu RH. Health-promoting components of fruits and vegetables in the diet. *Adv Nutr*. 2013;**4**(3):384S-92S. doi: [10.3945/an.112.003517](https://doi.org/10.3945/an.112.003517). [PubMed: [23674808](https://pubmed.ncbi.nlm.nih.gov/23674808/)].
38. Russo M, Spagnuolo C, Tedesco I, Bilotto S, Russo GL. The flavonoid quercetin in disease prevention and therapy: facts and fancies. *Biochem Pharmacol*. 2012;**83**(1):6-15. doi: [10.1016/j.bcp.2011.08.010](https://doi.org/10.1016/j.bcp.2011.08.010). [PubMed: [21856292](https://pubmed.ncbi.nlm.nih.gov/21856292/)].
39. Tamimi RM, Hankinson SE, Campos H, Spiegelman D, Zhang S, Colditz GA, et al. Plasma carotenoids, retinol, and tocopherols and risk of breast cancer. *Am J Epidemiol*. 2005;**161**(2):153-60. doi: [10.1093/aje/kwi030](https://doi.org/10.1093/aje/kwi030). [PubMed: [15632265](https://pubmed.ncbi.nlm.nih.gov/15632265/)].
40. Shen T, Khor SC, Zhou F, Duan T, Xu YY, Zheng YF, et al. Chemoprevention by lipid-soluble tea polyphenols in diethylnitrosamine/phenobarbital-induced hepatic pre-cancerous lesions. *Anticancer Res*. 2014;**34**(2):683-93. [PubMed: [24511000](https://pubmed.ncbi.nlm.nih.gov/24511000/)].
41. Crozier A, Clifford MN, Ashihara H. Plant secondary metabolites: occurrence, structure and role in the human diet. John Wiley & Sons; 2008.
42. W. Watson G, M. Beaver L, E. Williams D, H. Dashwood R, Ho E. Phytochemicals from cruciferous vegetables, epigenetics, and prostate cancer prevention. *AAPS J*. 2013;**15**(4):951-61. doi: [10.1208/s12248-013-9504-4](https://doi.org/10.1208/s12248-013-9504-4). [PubMed: [23800833](https://pubmed.ncbi.nlm.nih.gov/23800833/)].
43. Acharya A, Das I, Chandhok D, Saha T. Redox regulation in cancer: a double-edged sword with therapeutic potential. *Oxid Med Cell Longev*. 2010;**3**(1):23-34. doi: [10.4161/oxim.3.1.10095](https://doi.org/10.4161/oxim.3.1.10095). [PubMed: [20716925](https://pubmed.ncbi.nlm.nih.gov/20716925/)].
44. Levin DA. Plant Phenolics: An Ecological Perspective. *Am Nat*. 1971;**105**(942):157-81. doi: [10.1086/282712](https://doi.org/10.1086/282712).
45. Scalbert A, Williamson G. Dietary intake and bioavailability of polyphenols. *J Nutr*. 2000;**130**(8S Suppl):2073S-85S. [PubMed: [10917926](https://pubmed.ncbi.nlm.nih.gov/10917926/)].
46. Islam MB, Sarkar MMH, Shafique MZ, Jalil MA, Haque MZ, Amin R. Phytochemical screening and anti-microbial activity studies on lea macrophylla seed extracts. *J Sci Res*. 2013;**5**(2) doi: [10.3329/jsr.v5i2.13213](https://doi.org/10.3329/jsr.v5i2.13213).
47. Chirumbolo S. Plant phytochemicals as new potential drugs for immune disorders and cancer therapy: really a promising path?. *J Sci Food Agric*. 2012;**92**(8):1573-7. doi: [10.1002/jsfa.5670](https://doi.org/10.1002/jsfa.5670). [PubMed: [22473298](https://pubmed.ncbi.nlm.nih.gov/22473298/)].
48. Mateos R, Goya L, Bravo L. Uptake and metabolism of hydroxycinnamic acids (chlorogenic, caffeic, and ferulic acids) by HepG2 cells as a model of the human liver. *J Agric Food Chem*. 2006;**54**(23):8724-32. doi: [10.1021/jf061664g](https://doi.org/10.1021/jf061664g). [PubMed: [17090113](https://pubmed.ncbi.nlm.nih.gov/17090113/)].
49. Razzaghi-Asl N, Garrido J, Khazraei H, Borges F, Firuzi O. Antioxidant properties of hydroxycinnamic acids: a review of structure-activity relationships. *Curr Med Chem*. 2013;**20**(36):4436-50. [PubMed: [23834166](https://pubmed.ncbi.nlm.nih.gov/23834166/)].
50. Chaturvedi PK, Bhui K, Shukla Y. Lupeol: connotations for chemoprevention. *Cancer Lett*. 2008;**263**(1):1-13. doi: [10.1016/j.canlet.2008.01.047](https://doi.org/10.1016/j.canlet.2008.01.047). [PubMed: [18359153](https://pubmed.ncbi.nlm.nih.gov/18359153/)].
51. Gulcin I. Antioxidant activity of food constituents: an overview. *Arch Toxicol*. 2012;**86**(3):345-91. doi: [10.1007/s00204-011-0774-2](https://doi.org/10.1007/s00204-011-0774-2). [PubMed: [22102161](https://pubmed.ncbi.nlm.nih.gov/22102161/)].
52. Forster GM, Raina K, Kumar A, Kumar S, Agarwal R, Chen MH, et al. Rice varietal differences in bioactive bran components for inhibition of colorectal cancer cell growth. *Food Chem*. 2013;**141**(2):1545-52. doi: [10.1016/j.foodchem.2013.04.020](https://doi.org/10.1016/j.foodchem.2013.04.020). [PubMed: [23790950](https://pubmed.ncbi.nlm.nih.gov/23790950/)].
53. Murakami A. Modulation of protein quality control systems by food phytochemicals. *J Clin Biochem Nutr*. 2013;**52**(3):215-27. doi: [10.3164/jcbtn.12-126](https://doi.org/10.3164/jcbtn.12-126). [PubMed: [23704811](https://pubmed.ncbi.nlm.nih.gov/23704811/)].
54. Giardi MT, Touloupakis E, Bertolotto D, Mascetti G. Preventive or potential therapeutic value of nutraceuticals against ionizing radiation-induced oxidative stress in exposed subjects and frequent fliers. *Int J Mol Sci*. 2013;**14**(8):17168-92. doi: [10.3390/ijms140817168](https://doi.org/10.3390/ijms140817168). [PubMed: [23965979](https://pubmed.ncbi.nlm.nih.gov/23965979/)].
55. Raj Kapoor B, Muruges N, Chodon D, Sakthisekaran D. Chemoprevention of N-nitrosodiethylamine induced phenobarbital promoted liver tumors in rat by extract of Indigofera aspalathoides. *Biol Pharm Bull*. 2005;**28**(2):364-6. [PubMed: [15684501](https://pubmed.ncbi.nlm.nih.gov/15684501/)].
56. Stahl W, Sies H. Bioactivity and protective effects of natural carotenoids. *Biochim Biophys Acta*. 2005;**1740**(2):101-7. doi:

- 10.1016/j.bbadis.2004.12.006. [PubMed: 15949675].
57. Stahl W, Sies H. Antioxidant activity of carotenoids. *Mol Aspects Med.* 2003;**24**(6):345–51. [PubMed: 14585305].
58. Russo GL. Ins and outs of dietary phytochemicals in cancer chemoprevention. *Biochem Pharmacol.* 2007;**74**(4):533–44. doi: 10.1016/j.bcp.2007.02.014. [PubMed: 17382300].
59. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007;**39**(1):44–84. doi: 10.1016/j.biocel.2006.07.001. [PubMed: 16978905].
60. Deiters A, Martin SF. Synthesis of oxygen- and nitrogen-containing heterocycles by ring-closing metathesis. *Chem Rev.* 2004;**104**(5):2199–238. doi: 10.1021/cr0200872. [PubMed: 15137789].
61. Su C, Zhang P, Song X, Shi Q, Fu J, Xia X, et al. Tetrachlorobenzoquinone activates Nrf2 signaling by Keap1 cross-linking and ubiquitin translocation but not Keap1-Cullin3 complex dissociation. *Chem Res Toxicol.* 2015;**28**(4):765–74. doi: 10.1021/tx500513v. [PubMed: 25742418].
62. Chikamori K, Grozav AG, Kozuki T, Grabowski D, Ganapathi R, Ganapathi MK. DNA topoisomerase II enzymes as molecular targets for cancer chemotherapy. *Curr Cancer Drug Targets.* 2010;**10**(7):758–71. [PubMed: 20578986].
63. Bhandari PR. Crocus sativus L. (saffron) for cancer chemoprevention: A mini review. *J Tradit Complement Med.* 2015;**5**(2):81–7. doi: 10.1016/j.jtcme.2014.10.009. [PubMed: 26151016].
64. Ouyang L, Luo Y, Tian M, Zhang SY, Lu R, Wang JH, et al. Plant natural products: from traditional compounds to new emerging drugs in cancer therapy. *Cell Prolif.* 2014;**47**(6):506–15. doi: 10.1111/cpr.12143. [PubMed: 25377084].
65. Hemler ME. Tetraspanin proteins promote multiple cancer stages. *Nat Rev Cancer.* 2014;**14**(1):49–60. [PubMed: 24505619].
66. Sun B, Karin M. The therapeutic value of targeting inflammation in gastrointestinal cancers. *Trends Pharmacol Sci.* 2014;**35**(7):349–57. doi: 10.1016/j.tips.2014.04.011. [PubMed: 24881011].
67. Cichocki M, Dalek M, Szamalek M, Baer-Dubowska W. Naturally occurring phenolic acids modulate TPA-induced activation of EGFR, AP-1, and STATs in mouse epidermis. *Nutr Cancer.* 2014;**66**(2):308–14. doi: 10.1080/01635581.2014.864419. [PubMed: 24380573].
68. Shin JS, Yun CH, Cho YW, Baek NI, Choi MS, Jeong TS, et al. Indole-containing fractions of Brassica rapa inhibit inducible nitric oxide synthase and pro-inflammatory cytokine expression by inactivating nuclear factor-kappaB. *J Med Food.* 2011;**14**(12):1527–37. doi: 10.1089/jmf.2011.1611. [PubMed: 21877949].
69. Sun J, Chu YF, Wu X, Liu RH. Antioxidant and antiproliferative activities of common fruits. *J Agric Food Chem.* 2002;**50**(25):7449–54. [PubMed: 12452674].
70. Arumugam A, Stout M, Tsing C, Bhat A, Yong T, Nandy S, et al. Abstract 199: Short-term treatment with pregnancy levels of estradiol prevents breast cancer by delaying promotion and progression. *Cancer Res.* 2014;**73**(8 Supplement):199. doi: 10.1158/1538-7445.am2013-199.