Published online 2018 December 25.

**Research Article** 

# The Relationship Between Polymorphisms of Glutathione S-Transferase (GSTM1, GSTT1) Gene and Type 2 Diabetes Mellitus in Western Iran?

Fatemeh Kazemisafa <sup>1,\*</sup>, Kamalaldin Jadidian<sup>1,\*\*</sup>, Zohreh Rahimi<sup>2</sup>, Hamid Nomani<sup>2</sup>, Azadeh Ghanbari<sup>1</sup>, Gelareh Saeb<sup>1</sup> and Arash Alikhani<sup>1</sup>

<sup>1</sup>Health Department, Kermanshah University of Medical Sciences, Kermanshah, Iran

<sup>2</sup>Medical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

*corresponding author*: MS of Biochemistry, Health Department, Kermanshah University of Medical Sciences, Postal Code: 6718874414, Kermanshah, Iran. Tel: +98-8337257697, Email: kazemisafa.f@gmail.com

<sup>\*\*</sup> Corresponding author: MSc in Parasitology, Health Department, Kermanshah University of Medical Sciences, Postal Code: 6718874414, Kermanshah, Iran. Tel: +98-8337224043, Email: kamaljadid@yahoo.com

Received 2018 August 06; Revised 2018 December 05; Accepted 2018 December 11.

# Abstract

**Background:** Diabetes is a chronic metabolic disorder that causes many complications in the body. Oxidative free radicals affect human beta cells. Glutathione S-transferases (GSTs) can deactivate substances formed in oxidative stress.

**Objectives:** The present study was conducted to examine the relationship between GSTs polymorphisms and type 2 diabetes mellitus (T2DM), and to investigate the presence of GSTM1 and T1 genotypes in 176 T2DM patients and 174 healthy cases as controls. **Methods:** Multiplex PCR was used to confirm the presence or absence of GSTT1 and M1 genotypes.

**Conclusions:** The obtained results indicated that the polymorphism of GSTM1 and GSTT1 does not affect the prevalence of T2DM in the study population.

Keywords: Glutathione S-Transferase, Genetic Polymorphism, Oxidative Stress, T2DM

## 1. Background

Type 2 diabetes mellitus (T2DM) is the most prevalent type of diabetes that involves nearly 4% of the world population. This disease has a multifactorial origin that includes genetic and environmental agents (1).

This disease is characterized by metabolic irregularities causing long-term negative effects on the eyes, kidneys, nerves, and blood vessels (2).

Six percent of the world population is affected by T2DM, which is expected to rise to 77.6% by 2030(3, 4).

A total of 1.5 million diabetics have been reported in Iran(3, 5).

Increased levels of toxic substances and decreased antioxidant defense have been observed in T2DM (6-8). These situations can lead to oxidative stress with significant effects on diabetes prolongation (9-11).

Different genes are recognized to contribute to reducing the effect of toxic substances produced in oxidative stress (12-14), including one that produces Glutathione Stransferases (GSTs) as a multifunctional detoxification enzyme which defends cells against harmful effects caused by chemicals, metabolites and oxidative stress (15).

Several studies have identified GSTs polymorphisms as a risk factors for T2DM among different populations.

Given that oxidative stress plays a role in developing diabetes, GTS reduces oxidative stress in the body. The present study examined the association between this enzyme polymorphism (GSTT1 and GSTM1) and the prevalence of diabetes in the population of western Iran.

# 2. Methods

The present case-control study was performed in Kermanshah, Iran. The study participants comprised 176 patients, including 100 females and 76 males, aged 55.2  $\pm$  11.5 years, with T2DM confirmed by clinical tests, and 174 control volunteers, including 102 females and 72 males aged 55.3  $\pm$  11.2 years and proved by their medical history and laboratory tests not to have developed T2DM.

In the present study, the case group included T2DM patients with a fasting blood sugar (FBS) of at least 126 mg/dL

Copyright © 2018, Journal of Kermanshah University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited. and a 2hpp glucose of at least 200 mg/dL. Moreover, patients with type 1 diabetes, alcoholics, smokers and pregnant women were excluded from the study.

All the diabetics and controls were selected from the Regional Reference Laboratory, affiliated to Kermanshah University of Medical Sciences. Furthermore, the study was approved by the University Ethics Committee, and the participants submitted their consent forms prior to the commencement of the research.

Five mL of venous blood was taken and transferred to a test tube containing ethylene diamine tetra acetic acid (EDTA). Leukocytes were then isolated form the samples and their DNA was extracted.

The multiplex PCR technique helped detect the presence of GSTM1 and GSTT1 genes in the samples. Dihydrofolate Reductase (DHFR) locus was used as internal control.

The PCR primers included GSTM1 forward: 5' - GAA CTC CCT GAA AAGCTA AAGC-3', GSTM1 reverse: 5' GTT GGG CTC AAA TAT ACGGTG G-3' and GSTT1 forward: 5' - TTC CTT ACTGGT CCT CAC ATC TC-3', GSTT1 reverse: 5' - TCA CCGGA-CAT GGC CAG CA-3' and DHFR forward 5' -GGA ATG GAG AAC CAG GTC GTC TT-3', DHFR reverse 5'-GCA TGT CTT TGG GAT GTG GA-3'.

PCR amplification was performed when GSTM1 and GSTT1 were found to be present in the DNA genome of the case and control samples.

The presence or absence of GSTM1 (a 215-bp fragment) and GSTT1 (a 480-bp fragment) was approved in the presence of amplified fragments of the DHFR gene (280 bp) (Figure 1).



Figure 1. PCR products: Lines 1 and 4 indicate two GST1M1-positive samples. Line 6 indicates a GSTM1-positive sample. Lines 3 and 5 indicate two GSTT1-positive samples, and lines 2 and 7 show the elimination of GSTM1 and GSTT1.

# 3. Results

Table 1 presents the demographic characteristics of T2DM patients (n = 176) and the controls (n = 174). Accord-

ing to these results, the cases and controls showed no statistically significant differences in terms of age and gender (P > 0.05).

Fable 1. Characteristics of Patients and Controls							
Parameters		Patients	Controls	P Value			
Age, y		$55.2\pm11.5$	$55.3 \pm 11.2$	0.19			
Gender				0.46			
	Female, No. (%)	100 (56.8)	102 (58.6)				
	Male, No. (%)	76 (43.2)	72 (41.4)				
HbA1c (%)		$8.5\pm3.23$	$4.8\pm0.39$	< 0.001			
FBS, mg/dL		$167.5\pm36.9$	$86.38 \pm 7.7$	< 0.001			

Table 2 presents the prevalence of GSTM1 and GSTT1 in the cases and controls.

The prevalence of GSTM1-null was found to be 45.5% and that of GSTT1-null genotype to be 8.65% in the T2DM patients, whereas in the control group, the frequency of null genotypes of GSTM1 and GSTT1 was calculated as 48.3% and 17.75%, respectively.

Statistical analyses revealed that the polymorphism of GSTM1 (OR = 1.2; 95% CI 0.72 - 2.5, P = 0.6) and GSTT1 (OR = 1.3; 95% CI 0.91 - 2.11, P = 0.34) were not associated with the disease.

All the results obtained indicate that genotype distributions in the controls followed the Hardy-Weinberg equilibrium.

# 4. Discussion

T2DM is a harmful and globally prevalent disease. The disease complications exacerbate by different factors, including aging, prolonged obesity and decreased mobility (16).

Oxidative stress significantly affects the development of this disease (17). Given that  $\beta$ -cells carry a small amount of antioxidant agents, they are very susceptible to reactive oxygen species (ROS) damage. Ascented ROS can therefore play a key role in  $\beta$ -cell failures and the prolongation of insulin resistance and diabetes mellitus (17).

GST is an antioxidant enzyme that is responsible for protecting cells against oxidative stress by catalyzing the conjugation of glutathione to an extensive range of electrophiles (18).

According to the obtained results, no significant relationships were observed between GSTM1-null genotype and T2DM in the study population (45.5% vs. 48.3%).

No significant relationships were also observed between GSTT1-null genotype and T2DM in the study population (8.65% vs. 17.75%).

Genotypes	Patients, N = 100 <sup>a</sup>	Controls, N = 100 <sup>a</sup>	OR	95 % CI	P Value
GST M1					
Present	96 (54.5)	90 (51.7)	1.2	2.5 - 0.72	0.6
Null	80 (45.5)	84 (48.3)			
GSTT1					
Present	159 (91.32)	142 (82.5)	1.5	0.91 - 2.11	0.34
Null	17 (8.65)	32 (17.75)			

<sup>a</sup> Values are expressed as No. (%).

Several studies have been conducted in different populations, and different results have been obtained on the relationship of T2DM with the polymorphisms of GSTM1 and GSTT1.

In contrast to the present research, the study conducted by Moasser et al. (2) in southern Iran indicated that an increase in the GSTM1-null genotype can be associated with the pathogenesis of T2DM. Moreover, the studies conducted in Turkey and North India found GSTM1-null genotype to be a risk factor for developing T2DM (19, 20).

In line with the present findings, Chinese and Egyptian researchers (21, 22) found no significant differences between the GSTM1 genotype of the controls and the patients with T2DM; nevertheless their reports indicated that an increase in GSTT1-null is associated with developing T2DM.

A study by Amer et al. showed that the prevalence of GSTM1 and GSTT1 genotypes notably differs between T2DM patients and the controls in Egypt. Moreover, the combined genotype of GSTM1-null/GSTT1-null was found to increase the risk of developing T2DM (23).

A large body of research suggests that GSTs are essential for protecting DNA against harmful compounds that produce oxidative stress, and that the lack of these enzymes can cause susceptibility to many diseases (24), which has been proved to be associated with different environmental and genetic factors (25). Research suggest that different agents contribute to different levels of susceptibility to diseases in different populations, including physical activity, nutrition, stress, alcohol consumption, ethnic history and environmental conditions.

## 4.1. Conclusion

The present study showed that the frequency of GSTM1 and GSTT1 genotypes does not affect the prevalence of T2DM in the population of western Iran.

## Acknowledgments

The authors would like to express their gratitude to the employees of the Western Regional Reference Laboratory

for collaborating with this project.

#### Footnotes

**Conflict of Interests:** The authors declare that they have no conflict of interest to the publication of this article.

**Ethical Considerations:** The study was approved by the University Ethics Committee, and the participants submitted their consent forms prior to the commencement of the research.

**Funding/Support:** Kermanshah University of Medical Sciences, Kermanshah, Iran.

## References

- Horikawa Y, Oda N, Cox NJ, Li X, Orho-Melander M, Hara M, et al. Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. *Nat Genet*. 2000;**26**(2):163–75. doi: 10.1038/79876. [PubMed: 11017071].
- Moasser E, Kazemi-Nezhad SR, Saadat M, Azarpira N. Study of the association between glutathione S-transferase (GSTM1, GSTT1, GSTP1) polymorphisms with type II diabetes mellitus in southern of Iran. *Mol Biol Rep.* 2012;**39**(12):10187–92. doi: 10.1007/s11033-012-1893-4. [PubMed: 23014993].
- Azizi F, Guoya MM, Vazirian P, Dolatshati P, Habbibian S. Screening for type 2 diabetes in the Iranian national programme: A preliminary report. *East Mediterr Health J.* 2003;9(5-6):1122–7. [PubMed: 16450546].
- Hussain A, Vaaler S, Sayeed MA, Mahtab H, Ali SM, Khan AK. Type 2 diabetes and impaired fasting blood glucose in rural Bangladesh: A population-based study. *Eur J Public Health*. 2007;17(3):291–6. doi: 10.1093/eurpub/ckl235. [PubMed: 17008328].
- 5. Azizi F, Gouya MM, Vazirian P, Dolatshahi P, Habibian S. The diabetes prevention and control programme of the Islamic Republic of Iran. *East Mediterr Health J.* 2003;9(5-6):1114-21. [PubMed: 16450545].
- Giron MD, Salto R, Gonzalez Y, Giron JA, Nieto N, Periago JL, et al. Modulation of hepatic and intestinal glutathione S-transferases and other antioxidant enzymes by dietary lipids in streptozotocin diabetic rats. *Chemosphere*. 1999;**38**(13):3003–13. doi: 10.1016/S0045-6535(98)00502-5. [PubMed: 10230044].
- Opara EC. Oxidative stress, micronutrients, diabetes mellitus and its complications. J R Soc Promot Health. 2002;122(1):28–34. doi: 10.1177/146642400212200112. [PubMed: 11989140].
- 8. Fridlyand LE, Philipson LH. Oxidative reactive species in cell injury: Mechanisms in diabetes mellitus and therapeutic approaches. Ann

*NYAcad Sci.* 2005;**1066**:136–51. doi: 10.1196/annals.1363.019. [PubMed: 16533924].

- 9. West IC. Radicals and oxidative stress in diabetes. *Diabetic Medicine*. 2000;**17**(3):171-80. doi: 10.1046/j.1464-5491.2000.00259.x.
- Robertson RP, Harmon J, Tran PO, Tanaka Y, Takahashi H. Glucose toxicity in beta-cells: Type 2 diabetes, good radicals gone bad, and the glutathione connection. *Diabetes*. 2003;**52**(3):581-7. doi: 10.2337/diabetes.52.3.581. [PubMed: 12606496].
- Niedowicz DM, Daleke DL. The role of oxidative stress in diabetic complications. *Cell Biochem Biophys.* 2005;43(2):289–330. doi: 10.1385/CBB:43:2:289. [PubMed: 16049352].
- Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu Rev Pharmacol Toxicol.* 2005;45:51–88. doi: 10.1146/an-nurev.pharmtox.45.120403.095857. [PubMed: 15822171].
- Mates JM, Perez-Gomez C, Nunez de Castro I. Antioxidant enzymes and human diseases. *Clin Biochem.* 1999;**32**(8):595–603. doi: 10.1016/S0009-9120(99)00075-2. [PubMed: 10638941].
- Huang QY, Cheng MR, Ji SL. Linkage and association studies of the susceptibility genes for type 2 diabetes. *Acta Genetica Sinica*. 2006;**33**(7):573–89. doi:10.1016/s0379-4172(06)60087-5.
- Safa FK, Shahsavari G, Abyaneh RZ. Glutathione s-transferase M1 and T1 genetic polymorphisms in Iranian patients with glaucoma. *Iran J Basic Med Sci.* 2014;**17**(5):332–6. [PubMed: 24967061]. [PubMed Central: PMC4069838].
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004;**27**(5):1047-53. doi: 10.2337/diacare.27.5.1047. [PubMed: 15111519].
- Stephens JW, Khanolkar MP, Bain SC. The biological relevance and measurement of plasma markers of oxidative stress in diabetes and cardiovascular disease. *Atherosclerosis*. 2009;**202**(2):321–9. doi:

10.1016/j.atherosclerosis.2008.06.006. [PubMed: 18640680].

- Moussa SA. Oxidative stress in diabetes mellitus. *Romanian J biophys*. 2008;18(3):225–36.
- Yalin S, Hatungil R, Tamer L, Ates NA, Dogruer N, Yildirim H, et al. Glutathione S-transferase gene polymorphisms in Turkish patients with diabetes mellitus. *Cell Biochem Funct.* 2007;25(5):509–13. doi: 10.1002/cbf.1339. [PubMed: 16927413].
- Bid HK, Konwar R, Saxena M, Chaudhari P, Agrawal CG, Banerjee M. Association of glutathione S-transferase (GSTM1, T1 and P1) gene polymorphisms with type 2 diabetes mellitus in north Indian population. J Postgrad Med. 2010;56(3):176–81. doi: 10.4103/0022-3859.68633. [PubMed: 20739761].
- Wang G, Zhang L, Li Q. Genetic polymorphisms of GSTT1, GSTM1, and NQO1 genes and diabetes mellitus risk in Chinese population. *Biochem Biophys Res Commun.* 2006;**341**(2):310–3. doi: 10.1016/j.bbrc.2005.12.195. [PubMed: 16413497].
- 22. Rasool HAA, Kashmiry NK, Nowier SR, Morad H, Ismail S. Association of type 2 diabetes mellitus and glutathione S transferase (GSTM1 and GSTT1) genetic polymorphism. *Med J Cairo Univ*. 2009;**77**:71–7.
- Amer MA, Ghattas MH, Abo-Elmatty DM, Abou-El-Ela SH. Influence of glutathione S-transferase polymorphisms on type-2 diabetes mellitus risk. *Genet Mol Res.* 2011;10(4):3722–30. doi: 10.4238/2011.October.31.14. [PubMed: 22058002].
- Thier R, Bruning T, Roos PH, Rihs HP, Golka K, Ko Y, et al. Markers of genetic susceptibility in human environmental hygiene and toxicology: The role of selected CYP, NAT and GST genes. *Int J Hyg Environ Health.* 2003;206(3):149–71. doi: 10.1078/1438-4639-00209. [PubMed: 12872524].
- Prokopenko I, McCarthy MI, Lindgren CM. Type 2 diabetes: New genes, new understanding. *Trends Genet*. 2008;24(12):613–21. doi: 10.1016/j.tig.2008.09.004. [PubMed: 18952314].