

# A Comparison Between Job Stress and Insulin Resistance Among the Hospital Medical Staff

Parvin Zareian,<sup>1\*</sup> Fatemeh Ghasemi,<sup>1</sup> Seyd Reza Abtahi,<sup>2</sup> and Shahab Behzadi<sup>1</sup>

<sup>1</sup>Department of Physiology, AJA University of Medical Sciences, Tehran, IR Iran

<sup>2</sup>Department of Pharmacology, AJA University of Medical Sciences, Tehran, IR Iran

\*Corresponding author: Parvin Zareian, Department of Physiology, AJA University of Medical Sciences, Tehran, IR Iran. Tel/Fax: +98-2144609936, E-mail: zareianj2011@yahoo.com

Received 2016 March 16; Revised 2016 August 03; Accepted 2016 August 03.

## Abstract

**Background:** One of the important risk factors for insulin resistance is stress and a major source of stress is job stress.

**Objectives:** The current study aimed to determine the relationship between job stress score and insulin resistance among medical staff of the Imam Reza hospital, Tehran, Iran

**Methods:** The current descriptive cross-sectional study was performed on 97 medical staff of Imam Reza hospital, Tehran, Iran. The Osipow questionnaire was used to assess the level of occupational stress. Insulin resistance was calculated according to the formula: fasting insulin (microU/L)  $\times$  fasting glucose.

**Results:** There was no significant difference in insulin resistance among the hospital staff. Job stress was significantly higher in the intensive care unit staff than the staff of other departments ( $P < 0.01$ ). There was a significant positive correlation between insulin resistance and age, but there was no significant correlation between insulin resistance with body mass index (BMI) and stress score.

**Conclusions:** The intensive care unit (ICU) staff was exposed to more stress than the staff of other departments. But there was no relationship between job stress and insulin resistance among hospital staff. Further studies are needed to evaluate the relationship between job stress and insulin resistance.

**Keywords:** Stress, Insulin Resistance, Medical Staff

## 1. Background

Insulin resistance is a state in which cells are unable to respond to the normal actions of the insulin. Therefore, the blood sugar increases. High levels of blood sugar cause high levels of insulin production and it leads to down regulation of the insulin receptors. It means that there is a relative resistance to insulin despite the high level of insulin in the hospital staff. Previous studies showed that insulin resistance results in the cluster of lipid disorders, hypertension and glucose intolerance (1-4). Therefore, insulin resistance is a risk factor for developing heart disease and type 2 diabetes (5).

Several factors increase the risk of insulin resistance. Stress is one of the important risk factors. Stress results in various emotional and physical disorders including depression, anxiety, heart diseases, hypertension and immune system disturbances that increase vulnerability to infections (6-8). A major source of stress is job stress. Job stress is mentioned as a significant health problem. Tehrani et al., reported a significant relationship between job stress and mental health in a study on the effect of

stress on the nurses of emergency wards (9).

Stressful work situations such as: uncontrollable workloads, lack of job security, poor relationship with colleagues, lack of involvement in decision making, inadequate support from colleagues and superiors, dull and repetitive work, inadequate information about roles and responsibilities affecting both individual and organization lead to physiological responses that may finally contribute to sickness (10-12). Some reports show that medical staff faces high levels of stress at work because any medical errors or mistakes may threat patient's life which is sometimes irreversible. In addition, medical staff always faces suffering and death (13-15).

## 2. Objectives

The current study aimed to: 1, identify the job stress level among medical staff of the Imam Reza hospital; 2, measure the insulin resistance among medical staff of Imam Reza hospital; 3, determine the relationship between job stress score and insulin resistance.

### 3. Methods

#### 3.1. Participants

Participants included all the staff working in different departments of Imam Reza hospital, Tehran, Iran, from July to August 2013. Hospital departments included paraclinic, intensive care unit (ICU) and hospital wards.

Paraclinic department included odontology, pharmacy, pathology, radiology and central laboratory. Intensive care unit included ICU, coronary care unit (CCU) and emergency ward. Hospital wards included internal medicine, general surgery and operating rooms.

The Osipow questionnaire was used to assess the level of job stress among staff. The questionnaire consists of 60 statements based on a 5-point Likert scale; 5 for most of the times, 4 for usually, 3 for often, 2 for occasionally and 1 for never. Total score on this scale is considered in the assessment of occupational stress. Higher scores on this scale indicate more stress.

A copy of the occupational stress inventory (OSI) was given to the health care personnel. The completed questionnaires were analyzed by another person in the university.

Serum glucose level was measured by glucose oxidase-peroxidase (GOD-POD) method employing standard kits (Pars Azmoon, Karaj, Iran); fasting insulin level was also measured by enzyme linked immunosorbent assay (ELISA) using commercially available kits (Monobind, Inc. Lake Forest, CA, USA).

Insulin resistance was quantified using the homeostatic model assessment of insulin resistance (HOMA-IR) according to the formula:  $\text{fasting insulin (microU/L)} \times \text{fasting glucose (nm/L)} / 22.5$  as previously reported by Matthews et al. body mass index (BMI) is body weight in kilograms divided by the square of height in meters (16). All participants in this study signed the informed consent form and the study was approved by the ethical committee of the AJA University of Medical Sciences, Tehran, Iran.

#### 3.2. Statistical Analysis

All results were displayed as mean  $\pm$  SEM (standard error of mean). Statistical analysis of data was carried out by SPSS ver. 16. One-way analysis of variance and Post-Hoc tests were used to analyze different variables among the staff of different departments. Correlation analysis was performed to analyze the relationships between insulin resistance and dependent variables (age, job stress level and BMI).

Data analysis was performed using SPSS software version 16. P values  $\leq 0.05$  were considered significant.

### 4. Results

In total, 97 subjects (36 male and 61 female) participated in the study. Some anthropometric and clinical measurements of subjects are summarized in Table 1.

As indicated in Table 1, the staff of hospital wards was the youngest and those of ICU were the oldest ones. There was a significant difference between the staff of hospital wards and the ones in ICU regarding the age ( $P = 0.03$ ) (Figure 1). There was no significant difference among the staff of different departments regarding BMI.




Figure 1. The Comparison of Age Among the Staff of Different Departments of Imam Reza Hospital; \* $P < 0.05$

#### 4.1. Serum Insulin and Glucose Concentration

There was no significant difference regarding serum insulin and glucose concentration among the staff of different departments of the hospital (Figures 2 and 3).





Figure 2. The Comparison of Serum Glucose Concentration Among the Staff of Different Departments of Imam Reza Hospital

**Table 1.** Demographic and Laboratory Data of Study Subjects



Groups	Age, Y	Weight, kg	Height, m	BMI	Glucose, mg/dL	Insulin, $\mu$ IU/mL	Insulin Resistance	Stress Score
Paraclinical department staff (n = 40)	32.0 $\pm$ 1.1	66.6 $\pm$ 1.9	1.64 $\pm$ 0.01	25.0 $\pm$ 0.7	93.1 $\pm$ 2	91.32 $\pm$ 10.9	2.08 $\pm$ 0.2	162.1 $\pm$ 5.6
Intensive care unit staff (n = 21)	34 $\pm$ 1.7	71.3 $\pm$ 2.7	1.69 $\pm$ 0.02	24.9 $\pm$ 1.01	91 $\pm$ 3.1	89.5 $\pm$ 23	1.95 $\pm$ 0.47	175 $\pm$ 7.2
Hospital wards staff (n = 36)	30 $\pm$ 1	68.5 $\pm$ 2.1	1.67 $\pm$ 0.01	24.2 $\pm$ 0.4	90.5 $\pm$ 2.8	116.4 $\pm$ 27.1	2.47 $\pm$ 0.56	155.7 $\pm$ 4.8

Abbreviation: BMI, body mass index.

**Figure 3.** The Comparison of Serum Insulin Concentration Among the Staff of Different Departments of Imam Reza Hospital

#### 4.2. Insulin Resistance

According to [Figure 4](#), the staff of hospital wards and ICU showed the highest and lowest levels of insulin resistance respectively, but the difference between the rates was insignificant.

**Figure 4.** The Comparison of Serum Insulin Resistance Among the Staff of Different Departments of Imam Reza Hospital**Figure 5.** The Comparison of Job Stress Level Among the Staff of Different Departments of Imam Reza Hospital; \* $P < 0.05$ 

#### 4.3. Occupational Stress Level

Comparing occupational stress scores among hospital staff showed that the level of job stress among the staff of ICU was significantly higher than those of the hospital wards ( $P = 0.035$ ) ([Figure 5](#)).

#### 4.4. Correlations

There was a significant positive correlation between the insulin resistance and age, but there was no significant correlation between insulin resistance with BMI and stress score.

**Table 2.** Insulin Resistance

	Insulin Resistance	
	r	P
Age	0.3	0.004
BMI	0.01	0.9
Stress score	-0.09	0.3

Abbreviation: BMI, body mass index.

## 5. Discussion

Job stress adversely affects health at the individual level; it also negatively affects healthcare organizations and work output. At the individual level, stress is in association with metabolic syndrome and insulin resistance.

In the current study, there was no significant difference in insulin resistance among hospital departments, but there was a significant difference in age between the groups. Previous studies showed higher levels of insulin resistance in older people than younger ones and a positive relationship between age and insulin resistance (17, 18). Therefore, to study the effect of job stress on insulin resistance, age was considered as a confounding variable and ANCOVA was used to eliminate the effect of this factor. The results of ANCOVA as well as one-way ANOVA indicated no significant difference in insulin resistance between the three groups. Therefore, based on the results of the current study, insulin resistance does not depend on the working department, but it depends on the age.

However, in the current study, the relationship between age and insulin resistance was weak ( $r = 0.3$ ,  $P = 0.004$ ). The study by Karakelides et al. on the older adults showed that age-related reductions in insulin sensitivity was due to an age-related increase in adiposity rather than a consequence of advanced chronological age (19).

In addition, another study showed that insulin sensitivity in males up to 60 - 70 years was more depends on body fat mass than age (20).

A few published studies evaluated the source and level of stress among healthcare staff (15, 21); according to the finding of these studies, the main source of stress was heavy workload and inadequate staff. However, to the authors' best knowledge, the current study was the first study that compared insulin resistance and job stress level simultaneously among the staff of different hospital departments.

The results of the study showed that the staff of ICU, CCU and emergency ward experienced greater job stress than the other ones. This may not be surprising given that ICU and CCU staff usually faces high levels of stress at work, particularly due to the related clinical duties and death and that the ICU is especially stressful. Goodfellow et al. showed that ICU staff (physicians and nurses) exposed higher levels of stress than a normal working population (22).

In the current study, there was no correlation between insulin resistance and stress score. The study by Shiloah on 39 patients without diabetes and acute psychotic stress indicated an inverse correlation between  $\beta$ -cell function and insulin sensitivity with stress score (23). In another study, acute psychological stress (foot shock stress) in mice re-

sulted in the elevation of blood glucose levels in both glucose tolerance and insulin tolerance tests (24). The difference may be due to the type of stress and the method used to estimate insulin resistance.

To assess insulin resistance in the subjects, hyperinsulinemic euglycemic clamp is the best technique, but it was costly, time-consuming and required more cooperation of hospital staff. Consequently, this method did not use, which was a limitation of the present study.

In spite of the current study, some previous studies reported a correlation between insulin resistance and BMI (25, 26). Trirogoff et al. reported that the primary determinant of insulin resistance in patients with chronic kidney disease (CKD) is BMI, specifically, fat mass. Another study on 132 patients with type 2 diabetes showed as BMI increased, the patients exhibited higher levels of insulin resistance and lower levels of insulin sensitivity. In this study glucose tolerance test was used to assess insulin resistance and insulin sensitivity indexes (25).

A longitudinal population-based cohort study showed that one standard deviation increase in rate of change in BMI corresponded to a decrease in insulin sensitivity by 19%; this study also quantified insulin sensitivity using a hyperinsulinemic euglycemic clamp (27).

The basis of this difference may be due to the study population and various methods used to estimate insulin resistance.

The current study had several limitations. First, there were small number of subjects in some departments and the cooperation of physicians was poor.

Low number of participants in the study results in low power of statistical analysis. The insulin resistance did not measure using the hyperinsulinemic euglycemic clamp. As previously mentioned, this is the best method to evaluate insulin resistance.

### 5.1. Conclusion

It seems that the staff of ICU is exposed to more job stress than other department staff, but there was no relationship between the job stress and insulin resistance among hospital staff. In addition, it is better to pay more attention to the age of the selected subjects working in CCU, ICU and emergency ward. However, further studies are needed to test job stress and insulin resistance in the other hospitals affiliated to AJA University to assess the correlation between job stress score and insulin resistance level.

### Acknowledgments

Authors are grateful to Dr. Mohammad Reza Ebrahimi, department of psychology, school of medicine, AJA University of Medical Sciences, Tehran, Iran, for providing Osipow

questionnaire to measure the level of occupational stress. This research was granted by AJA University of Medical Sciences (code no. 992241). Authors wish to thank the staff of Imam Reza Hospital affiliated to AJA University of Medical Sciences for their participation in the study.

## Footnotes

**Authors' Contribution:** Study concept and design, Parvin Zareian; acquisition of data, Shahab Behzadi and Fatemeh Ghasemi; analysis and interpretation of data, Parvin Zareian and Seyd Reza Abtahi; drafting of the manuscript, Parvin Zareian; critical revision of the manuscript for important intellectual content, Parvin Zareian; statistical analysis, Parvin Zareian and Seyd Reza Abtahi; administrative, technical and material support, Imam Reza hospital and Mohammad Reza Ebrahemi; study supervision: Parvin Zareian

**Financial Disclosure:** There was no financial disclosure regarding the material in the manuscript.

## References

1. Reaven GM. Role of Insulin Resistance in Human Disease. *Diabetes*. 1988;**37**(12):1595–607. doi: [10.2337/diab.37.12.1595](https://doi.org/10.2337/diab.37.12.1595).
2. DeFronzo RA, Ferrannini E. Insulin resistance. A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diabetes Care*. 1991;**14**(3):173–94. [PubMed: [2044434](https://pubmed.ncbi.nlm.nih.gov/2044434/)].
3. Cambien F, Warnet JM, Eschwege E, Jacqueson A, Richard JL, Rosselin G. Body mass, blood pressure, glucose, and lipids. Does plasma insulin explain their relationships?. *Arteriosclerosis*. 1987;**7**(2):197–202. [PubMed: [3555432](https://pubmed.ncbi.nlm.nih.gov/3555432/)].
4. Ferrannini E, Haffner SM, Mitchell BD, Stern MP. Hyperinsulinaemia: the key feature of a cardiovascular and metabolic syndrome. *Diabetologia*. 1991;**34**(6):416–22. [PubMed: [1884900](https://pubmed.ncbi.nlm.nih.gov/1884900/)].
5. Groop L, Forsblom C, Lehtovirta M, Tuomi T, Karanko S, Nissen M, et al. Metabolic consequences of a family history of NIDDM (the Botnia study): evidence for sex-specific parental effects. *Diabetes*. 1996;**45**(11):1585–93. [PubMed: [8866565](https://pubmed.ncbi.nlm.nih.gov/8866565/)].
6. Bunker SJ, Colquhoun DM, Esler MD, Hickie IB, Hunt D, Jelinek VM, et al. "Stress" and coronary heart disease: psychosocial risk factors. *Med J Aust*. 2003;**178**(6):272–6. [PubMed: [12633484](https://pubmed.ncbi.nlm.nih.gov/12633484/)].
7. Mayer EA. The neurobiology of stress and gastrointestinal disease. *Gut*. 2000;**47**(6):861–9. [PubMed: [11076888](https://pubmed.ncbi.nlm.nih.gov/11076888/)].
8. O'Leary A. Stress, emotion, and human immune function. *Psychol Bull*. 1990;**108**(3):363–82. [PubMed: [2270233](https://pubmed.ncbi.nlm.nih.gov/2270233/)].
9. Tehrani H, Rakhshani T, Shojaee Zadeh D, Hosseini SM, Bagheriyan S. Analyzing the relationship between job stress to mental health, personality type and stressful life events of the nurses occupied in tehran 115 emergency. *Iran Red Crescent Med J*. 2013;**15**(3):272–3. doi: [10.5812/ircmj.1917](https://doi.org/10.5812/ircmj.1917). [PubMed: [23984013](https://pubmed.ncbi.nlm.nih.gov/23984013/)].
10. Kivimaki M, Nyberg ST, Batty GD, Fransson EI, Heikkila K, Alfredsson L, et al. Job strain as a risk factor for coronary heart disease: a collaborative meta-analysis of individual participant data. *Lancet*. 2012;**380**(9852):1491–7. doi: [10.1016/S0140-6736\(12\)60994-5](https://doi.org/10.1016/S0140-6736(12)60994-5). [PubMed: [22981903](https://pubmed.ncbi.nlm.nih.gov/22981903/)].
11. Boscolo P. [Effects of occupational stress and job insecurity on the immune response]. *G Ital Med Lav Ergon*. 2009;**31**(3):277–80. [PubMed: [19943442](https://pubmed.ncbi.nlm.nih.gov/19943442/)].
12. Huerta-Franco MR, Vargas-Luna M, Tienda P, Delgado-Holtfort I, Balleza-Ordaz M, Flores-Hernandez C. Effects of occupational stress on the gastrointestinal tract. *World J Gastrointest Pathophysiol*. 2013;**4**(4):108–18. doi: [10.4291/wjgp.v4.i4.108](https://doi.org/10.4291/wjgp.v4.i4.108). [PubMed: [24244879](https://pubmed.ncbi.nlm.nih.gov/24244879/)].
13. Firth-Cozens J, Greenhalgh J. Doctors' perceptions of the links between stress and lowered clinical care. *Soc Sci Med*. 1997;**44**(7):1017–22. [PubMed: [9089922](https://pubmed.ncbi.nlm.nih.gov/9089922/)].
14. Cydulka RK, Emerman CL, Shade B, Kubincanek J. Stress levels in EMS personnel: a national survey. *Prehosp Disaster Med*. 1997;**12**(2):136–40. [PubMed: [10186997](https://pubmed.ncbi.nlm.nih.gov/10186997/)].
15. Escriba-Aguir V, Bernabe-Munoz Y. Job stress perceived by hospital medical staff [in Spanish]. *Gac Sanit*. 2002;**16**(6):487–96. [PubMed: [12459131](https://pubmed.ncbi.nlm.nih.gov/12459131/)].
16. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;**28**(7):412–9. [PubMed: [3899825](https://pubmed.ncbi.nlm.nih.gov/3899825/)].
17. Refaie MR, Sayed-Ahmed NA, Bakr AM, Aziz MYA, El Kannishi MH, Abdel-Gawad SS. Aging is an Inevitable Risk Factor for Insulin Resistance. *J Taibah Univ Med Sci*. 2006;**1**(1):30–41.
18. Fink RI, Kolterman OG, Griffin J, Olefsky JM. Mechanisms of insulin resistance in aging. *J Clin Invest*. 1983;**71**(6):1523–35. [PubMed: [6345584](https://pubmed.ncbi.nlm.nih.gov/6345584/)].
19. Karakelides H, Irving BA, Short KR, O'Brien P, Nair KS. Age, obesity, and sex effects on insulin sensitivity and skeletal muscle mitochondrial function. *Diabetes*. 2010;**59**(1):89–97. doi: [10.2337/db09-0591](https://doi.org/10.2337/db09-0591). [PubMed: [19833885](https://pubmed.ncbi.nlm.nih.gov/19833885/)].
20. Boden G, Chen X, DeSantis RA, Kendrick Z. Effects of age and body fat on insulin resistance in healthy men. *Diabetes Care*. 1993;**16**(5):728–33. [PubMed: [8495612](https://pubmed.ncbi.nlm.nih.gov/8495612/)].
21. Saha D, Sinha R, Bhavsar K. Understanding job stress among health-care staff. *J Health and Allied Sci*. 2011;**10**(1).
22. Goodfellow A, Varnam R, Rees D, Shelly MP. Staff stress on the intensive care unit: a comparison of doctors and nurses. *Anaesthesia*. 1997;**52**(11):1037–41. [PubMed: [9404163](https://pubmed.ncbi.nlm.nih.gov/9404163/)].
23. Shiloah E, Witz S, Abramovitch Y, Cohen O, Buchs A, Ramot Y, et al. Effect of acute psychotic stress in nondiabetic subjects on beta-cell function and insulin sensitivity. *Diabetes Care*. 2003;**26**(5):1462–7. [PubMed: [12716805](https://pubmed.ncbi.nlm.nih.gov/12716805/)].
24. Li L, Li X, Zhou W, Messina JL. Acute psychological stress results in the rapid development of insulin resistance. *J Endocrinol*. 2013;**217**(2):175–84. doi: [10.1530/JOE-12-0559](https://doi.org/10.1530/JOE-12-0559). [PubMed: [23444388](https://pubmed.ncbi.nlm.nih.gov/23444388/)].
25. Tirirogoff ML, Shintani A, Himmelfarb J, Ikizler TA. Body mass index and fat mass are the primary correlates of insulin resistance in non-diabetic stage 3–4 chronic kidney disease patients. *Am J Clin Nutr*. 2007;**86**(6):1642–8. [PubMed: [18065581](https://pubmed.ncbi.nlm.nih.gov/18065581/)].
26. Chung JO, Cho DH, Chung DJ, Chung MY. Associations among body mass index, insulin resistance, and pancreatic beta-cell function in Korean patients with new-onset type 2 diabetes. *Korean J Intern Med*. 2012;**27**(1):66–71. doi: [10.3904/kjim.2012.27.1.66](https://doi.org/10.3904/kjim.2012.27.1.66). [PubMed: [22403502](https://pubmed.ncbi.nlm.nih.gov/22403502/)].
27. Riserus U, Arnlov J, Berglund L. Long-term predictors of insulin resistance: role of lifestyle and metabolic factors in middle-aged men. *Diabetes Care*. 2007;**30**(11):2928–33. doi: [10.2337/dc07-0360](https://doi.org/10.2337/dc07-0360). [PubMed: [17644620](https://pubmed.ncbi.nlm.nih.gov/17644620/)].