



Protective Effect of Thymoquinone on D-Galactose-Induced Aging in Mice

Hasan Badie Bostan¹, Soghra Mehri², Elaheh Mohammadi¹ and Hossein Hosseinzadeh^{2,*}

¹Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

²Department of Pharmacodynamics and Toxicology, Pharmaceutical Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Department of Pharmacodynamics and Toxicology, Pharmaceutical Research Center, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. Tel: +98-5138819042, Fax: +98-5138823251, Email: hosseinzadehh@mums.ac.ir

Received 2017 May 23; Accepted 2018 May 30.

Abstract

Background: Aging is considered as a gradual biological process, which induces alteration in biochemistry and morphology of a biological system. Thymoquinone (TQ), is the major ingredient of volatile oil from *Nigella sativa* L. seeds. Because of the various pharmacological effects of TQ, this constituent was used for the current study.

Objectives: Anti-aging property of TQ in D-galactose-induced model of aging was investigated in mice.

Methods: Aging was induced in Razi mice with D-galactose injection (500 mg/kg, SC, daily) for 42 consecutive days. Animal treatment conducted with D-galactose alone or with different doses of TQ (2.5, 5 and 10 mg/kg). Malondialdehyde (MDA) and glutathione (GSH) concentration were determined in liver and brain tissues at the end of treatment. Serum concentration of IL-6 and tumor necrosis factor- α (TNF- α), alanine aminotransferase (ALT) and aspartate aminotransferase (AST) as well as dehydroepiandrosterone-sulfate (DHEA-S) and testosterone were determined.

Results: Administration of D-galactose (500 mg/kg, SC, daily) for 42 consecutive days increased the level of MDA in brain and liver tissues while diminished the GSH content. Additionally, the serum levels of TNF- α , IL-6, ALT and AST were significantly elevated following induction of aging phenomena while the level of male sex hormones were markedly reduced. Treatment with TQ (5 and 10 mg/kg) elevated level of GSH while decreased MDA production. In addition, the serum level of TNF- α , IL-6, ALT and AST decreased while the level of male sex hormones improved following treatment with TQ.

Conclusions: TQ showed anti-aging effect in this model. These effects in part can be mediated through antioxidant effects of TQ.

Keywords: Thymoquinone, *Nigella sativa*, Oxidative Stress, Cytokines, Antioxidant

1. Background

Since many years ago, people have been found herbal products to be promising agents for treatment of disease. Many natural medicines have low adverse effects, price and good efficacy in different human diseases (1). Anti-aging features of some herbs which are used traditionally have been mentioned in new experimental clinical studies (2).

Nigella sativa L. which is also known as black seed belongs to the Ranunculaceae family. Remarkable biological properties of *N. sativa* seeds is attributed to thymoquinone (TQ). Up to now, it was applied for various diseases and conditions such as gastrointestinal problems, asthma, high blood pressure, diabetes, inflammatory conditions, fever, infection (3-5) and convulsion (6).

From the pharmacological point of view, TQ is a quinone showing anti-inflammatory and analgesic effects (7), protective effects against chemical induced carcino-

genesis (8), and also is a powerful inhibitor of eicosanoid production, especially thromboxane B₂ and leukotriene B₄ (9). TQ has more inhibitory effect on PGE₂ production than indomethacin (10). Because of antioxidant effect, TQ is able to protect hepatocyte membranes from peroxidation (11, 12).

In a model of hippocampus injury following transient forebrain ischemia in rats, TQ showed protective effects. It was concluded that TQ elevated GSH content, SOD and catalase activity (13). TQ markedly inhibited A β 1-40 induced apoptosis in cerebellar granule neurons mainly via intrinsic/extrinsic caspase cascades (14).

Aging is a complex biological event which demonstrated by induction of harmful changes during the time. Oxidative stress theory of aging and free radical generation have been mentioned as the most important interpretation for the process of aging (15). Deregulations of immune system followed by elevation of proinflammatory

cytokines, reduction of sex hormone levels and induction of lipid peroxidation are different factors which can be changed during this process (16). A typical model of aging using D-galactose was selected in the present study.

D-galactose as a reducing sugar can be metabolized at normal concentrations. Aldose and hydroperoxidase are by-products of D-galactose, which induce the production of superoxide anion and oxygen-originated free radicals (17).

2. Objectives

In regard to potent antioxidant properties of TQ, this study was conducted to examine the anti-aging potential effect of TQ on D-galactose induced aging model in mice using GSH and MDA concentration, sex hormone and inflammatory indices determination.

3. Methods

3.1. Chemicals

D-galactose, TQ, malondialdehyde tetrabutylammonium and DTNB (5, 5'-dithiobis-(2-nitrobenzoic acid)) were obtained from Sigma. *TNF- α* ELISA kit and *IL-6* ELISA kit were obtained from eBioscience.

3.2. Animal

Male mice, 25 - 30 g were housed in colony rooms with ($21 \pm 2^\circ\text{C}$) under a 12/12 h light/dark cycle with free access to food and water. All experiments were performed according to Mashhad University of Medical Sciences, Ethical Committee Acts (920979).

All animals for this study were provided from Faculty of Pharmacy, Mashhad University of Medical Sciences.

3.3. Experimental Design

Induction of aging by D-galactose has been considered as a valuable model of aging in different studies (2, 18, 19). Therefore in the current experiment, aging was induced following administration of D-galactose (500 mg/kg, SC, daily) for 42 days to mice (2).

For our study, animals were divided into 6 groups (n = 10) randomly and treatment was done as described below:

- 1- Control, normal saline, SC.
- 2- D-galactose (500mg/kg) SC for 42 days.
- 3- D-galactose (500mg/kg) SC for 42 days + TQ (2.5 mg/kg) intraperitoneally (IP) (11).
- 4- D-galactose (500mg/kg) SC for 42 days + TQ (5 mg/kg) IP (11).
- 5- D-galactose (500mg/kg) SC for 42 days + TQ (10 mg/kg) IP (11).

6- TQ (10 mg/kg) IP.

TQ was administrated daily to animals intraperitoneally. After the treatment period, cardiac blood samples were collected under anesthesia. Serum samples were taken by centrifuging blood at 4°C for 10 minutes and were transferred to microtubes for separated biochemical assay. By means of liquid nitrogen, separated liver and brain tissues were snap-frozen and kept in -80 freezers until analyzing day.

3.4. Lipid Peroxidation Assay

Using malondialdehyde (MDA) content determination, the level of lipid peroxidation was assessed. The reaction of MDA with thiobarbituric acid (TBA) results in a complex with pink color which can be detected in λ_{max} of 532 nm.

At the first step, brain or liver tissue homogenate 10% in KCl was prepared. Then 3 mL phosphoric acid (1%), 1 mL TBA (0.6%) and 0.5 mL of tissue homogenate were mixed. After that, the mixtures were heated in a boiling water bath for 45 minutes. N-butanol was added to the mixtures and vortex-mixed for 1 minute. The mixtures were centrifuged at 3000 g for 10 minutes. Finally, the absorbance of organic layers were measured at 532 nm (20). Malondialdehyde tetrabutyl ammonium was used to plot calibration curve. MDA quantity was reported as nmol/g tissue.

3.5. Reduced Glutathione (GSH) Assay

With some modification, samples were analyzed for determination of brain and liver GSH content by use of previously published method (Moron et al.). For this method, homogenized samples were mixed with 10% tricolor acetic acid (TCA) [600 μL , 1:1 (v:v)] and vortexed. The supernatants of these mixtures were obtained by centrifuge at 2500 g for 10 minutes. Then the supernatants were separated and mixed with reaction mixtures including 2 mL phosphate buffer (pH: 8) and 500 μL DTNB indicator [5,5' di thiobis-(2-nitrobenzoic acid)]. The absorbance of final complex was recorded at 412 nm (21) using a spectrophotometer (Jenway 6105 uv/vis, UK). Commercially available GSH was used to plot calibration curve. GSH Levels were reported as nmol/g tissue.

3.6. Measurement of *TNF- α* and *IL-6*

Determination of *TNF- α* and *IL-6* concentrations in serum conducted by *TNF- α* ELISA kit and *IL-6* ELISA kit (eBioscience). The absorbance of the final colored complex was calculated in 450 nm as the primary wave length and 650 nm as the reference wavelength. *TNF- α* and *IL-6* levels were expressed as pg/mL.

3.7. Measurement of Testosterone and dehydroepiandrosterone - Sulfate (DHEA-S)

Levels of these sex hormones were determined in accordance with kit protocols. Data were expressed as ng/mL (testosterone) and ng/dL (DHEA-S).

3.8. Measurement of ALT and AST (IU/L)

These markers were measured by Mindary autos Analyzer (BS 800) according to kit protocols. Data were expressed as IU/L.

3.9. Statistical Analysis

Statistical analysis was performed using one-way ANOVA followed by Tukey post-hoc test for multiple comparisons to evaluate the discrepancy of the data. P-value less than 0.05 were considered statistically significant.

4. Results

4.1. Effect of TQ on Lipid Peroxidation Induced by D-Galactose in Mice

As shown in [Figure 1A](#), MDA level in D-galactose treated group was significantly higher than control one, ($P < 0.05$). Treatment with 2.5, 5 and 10 mg/kg of TQ reduced lipid peroxidation in brain tissue, when compared to aged mice ($P < 0.05$).

In addition, exposure to D-galactose for 42 days significantly increased lipid peroxidation in liver tissue in comparison to control group ($P < 0.01$). Interestingly, TQ (2.5, 5 and 10 mg/kg) markedly inhibited lipid peroxidation when compared to D-galactose treated animals ($P < 0.001$). Results have been shown in [Figure 1B](#).

TQ (10 mg/kg) alone did not induce lipid peroxidation in both brain and liver tissues in comparison with control group.

4.2. Effect of TQ on Tissue GSH Content in Aged Mice

As presented in [Figure 2A and B](#), TQ administration (5 and 10 mg/kg) significantly increased the reduced content of GSH in brain and liver of aged animals, ($P < 0.05$).

Administration of TQ (10 mg/kg) alone did not change GSH content when compared to control group.

4.3. Effect of TQ on ALT and AST Levels in Serum of Aged Mice

AST and ALT levels markedly increased in D-galactose-received group. In comparison with D-galactose aged mice, treatment with TQ (5 mg/kg) diminished level of AST and ALT ($P < 0.05$ for AST and $P < 0.01$ for ALT). As depicted in [table 1](#), no remarkable change was detected in level of serum ALT and AST following treatment with TQ (10 mg/kg) alone when compared to control animals.

4.4. Effect of TQ on Sex Hormone Levels of Aged Mice

Sex hormones as another marker that can be changed following aging were determined using Elisa kit. Treatment of animals with D-galactose reduced level of testosterone from 15 ± 2.68 ng/dL in control animals to 6 ± 3.22 ng/mL in aged ones. Also, the level of DHEA-S was 40 ± 1.63 ng/mL in control group that reduced to 26 ± 9.29 ng/mL in aged animals. TQ (2.5 and 5 mg/kg) significantly recovered both levels of testosterone and DHEA-S in aged animals ([Table 1](#)).

4.5. Effect of TQ on Level of TNF- α and IL-6 in D-Galactose-Received Mice

As presented in [Figure 3A and B](#), TNF- α and IL-6 levels in aged mice were higher than control group.

As shown in [Figure 3A and B](#), the level of TNF- α and IL-6 were considerably higher in D-galactose-received mice in comparison to control group ($P < 0.001$), while administration of TQ (5 and 10 mg/kg) significantly reversed the mentioned inflammatory markers. No significant change was observed in the level of IL-6 and TNF- α following exposure to TQ (10 mg/kg) alone.

5. Discussion

The anti-aging effect of TQ in model of aging induced with D-galactose (SC, for 42 days) in mice was illustrated in this study. Aging process is known by significant oxidative stress that is observed with reduction of GSH content in brain and liver tissues and consequently elevation of MDA level (marker of lipid peroxidation). Biochemical factors including ALT, AST, IL-6 and TNF- α increased following the aging process. TQ with doses of 5 and 10 mg/kg significantly inhibited brain and liver injury, reduced inflammation and increased level of sex hormones.

Importance of oxidative stress in aging pathogenesis is undeniable. It has been exhibited that aging is associated with a remarkable reduction in mitochondrial function, decline of GSH, enhancement of free radical production and progression of lipid peroxidation (2).

It should be mentioned that liver injury can be observed following aging (22). Change in MDA and GSH levels in aged animals of our study was in accordance with other works. Also, ALT and AST concentrations were remarkably elevated in aged animals in comparison with control group. Interestingly, treatment of animals with TQ (5 and 10 mg/kg) could reduce liver injury which was determined with reduction of serum enzymes, elevation of GSH content and finally diminishing lipid peroxidation.

Hepatoprotective effects of TQ have been evaluated in different studies. For instance, administration of TQ to




Figure 1. Effect of TQ on lipid peroxidation induced by D-galactose in brain (A) and liver (B). Data are expressed as mean \pm SD (n=7). # P < 0.05, ## P < 0.01 vs. control, *P < 0.05, *** P < 0.001 vs. D-galactose treated animals. DG (D-galactose), TQ (thymoquinone), CON (control).




Figure 2. Effect of TQ on brain (A) and liver (B) GSH content following treatment of animals with D-galactose. Data are expressed as mean \pm SD (n=7). ## P < 0.01 and ### P < 0.001 vs. control, *P < 0.05 and ** P < 0.01 vs. D-galactose treated animals. DG (D-galactose), TQ (thymoquinone), CON (control).

Sprague-Dawley rats reduced MDA content in liver, which means inhibition of lipid peroxide generation (23). TQ reduced the level of serum enzymes, increased the hepatic GSH content and consequently recovered hepatotoxicity of carbon tetrachloride (24).

There is a strong association between oxidative stress and neurodegenerative diseases (25).

Nowadays, induction of aging with D-galactose is well

accepted to provide a model of neurotoxicity, memory dysfunction, and oxidative damage (26). We observed that D-galactose administration to mice could induce brain damage, while TQ significantly inhibited lipid peroxidation in aged animals.

TQ exhibited neuroprotective effects in a model of hippocampus injury induced by transient forebrain ischemia. Enhancement of GSH level and CAT and SOD activity have

Table 1. Effect of TQ on ALT, AST and Sex Hormones Levels in Serum of D-Galactose-Induced Aged Mice^a

Test	AST (IU/L)	ALT (IU/L)	Testosterone (ng/dL)	DHEA-S (ng/mL)
Control	246.4 ± 7.57	120.3 ± 7.09	15 ± 2.68	40 ± 1.63
D-galactose (DG)	335 ± 32.66 ^b	176.37 ± 14.59 ^b	6 ± 3.22 ^b	26 ± 9.29 ^c
DG+TQ 2.5 mg/kg	256.6 ± 57.12 ^d	134.5 ± 30.28	13.1 ± 2.5 ^d	36.85 ± 4.74 ^d
DG+TQ 5 mg/kg	259.42 ± 31.96 ^d	109.75 ± 37.76 ^c	13.66 ± 3.7 ^d	36.28 ± 5.4 ^d
DG+TQ 10 mg/kg	289.57 ± 48.8	175.57 ± 57.34	6.4 ± 4.24	26.4 ± 2.45
TQ 10 mg/kg	249 ± 29.48	110.3 ± 17.53	17.8 ± .616	35.71 ± 8.44

Abbreviations: DG, D-galactose; TQ, thymoquinone; CON, control; AST, aspartate aminotransferase; ALT, alanine amino transferase; DHEA-S, dehydroepiandrosterone-sulfate.

^a Values are expressed as mean ± SD (n = 7).

^b P < 0.05 vs. control.

^c P < 0.01 vs. control.

^d P < 0.05 vs. D-galactose treated animals.

^e P < 0.01 vs. D-galactose treated animals.




Figure 3. Effect of TQ on *TNF-α* (A) and *IL-6* (B) levels in serum of D-galactose-induced aged mice. Data are expressed as mean ± SD (n = 5). ### P < 0.001 vs. control, *P < 0.01, **P < 0.05 and ***P < 0.001 vs. D-galactose treated animals. DG (D-galactose), TQ (thymoquinone), CON (control).

been revealed in TQ-neuroprotective mechanisms. TQ markedly reduced A β 1-40 induced apoptosis in cerebellar granule neurons (14). In cerebral cortex of Wistar rats, TQ decreased peroxidation of lipid induced with acrylamide (27).

Aging process is related with a disturbance in immune system function, induction of inflammation and elevation of proinflammatory cytokines (e.g., *IL-6*, *TNF-α*). While the data are not uniform, *IL-6* and *TNF-α* are thought to be involved in morbidity and mortality in the elderly, and *IL-6* has been regarded as the strongest risk marker in healthy elderly people (28). Studies indicated that, in healthy old populations, elevation in circulating *IL-6* represents a systemic response to local pro-inflammatory action. Long

term elevated *TNF-α* and *IL-6* have several biological activities that induce age-associated pathology and mortality (28).

In the current study level of these important cytokines elevated in aged animals. Interestingly, TQ (2.5, 5 and 10 mg/kg) could reduce the level of increased cytokines.

In the inflammation cascade, TQ plays a protective role with inhibition of production of leukotriene B₄ and thromboxane B₂ in eicosanoid biosynthesis pathway (9).

Changes of the level of sex hormones have been considered as an important marker during aging (29). Also, there is a correlation between inflammatory markers and testosterone concentration (30). In regard to these facts, we decided to evaluate serum testosterone and DHEA-S in aged

mice. In agreement to recent reports, the level of these factors markedly diminished in aged animals in comparison to control, but treatment with TQ (5 and 10 mg/kg) significantly increased sex hormone levels. A significant elevation in testosterone concentration in rats was observed following oral administration of water extract of *N. sativa* (31).

5.1. Conclusions

The present study clearly indicated that TQ possess significant anti-aging effects in the model of aging induced by D-galactose especially through enhancement of GSH content, decline of lipid peroxidation and inhibition of inflammation.

Acknowledgments

Authors are thankful to the vice chancellor of research, Mashhad University of Medical Sciences for financial support. The results explained in this paper were part of a Master of Science degree thesis.

Footnotes

Authors' Contribution: Hossein Hosseinzadeh: study concept, design and critical revision of the manuscript for important intellectual content; Soghra Mehri: drafting of the manuscript and advisor; Hasan Badie Bostan: doing experiments and drafting of the manuscript; Elaheh Mohammadi: doing experiments.

Conflict of Interests: The authors declare not to have any conflict of interests.

Ethical Considerations: All experiments were performed according to Mashhad University of Medical Sciences, Ethical Committee Acts (920979).

Funding/Support: This research was supported by a grant of vice chancellor of research from Mashhad University of Medical Sciences (920979).

References

- Hasani-Ranjbar S, Larijani B, Abdollahi M. A systematic review of the potential herbal sources of future drugs effective in oxidant-related diseases. *Inflamm Allergy Drug Targets*. 2009;**8**(1):2-10. doi: [10.2174/18752809787582561](https://doi.org/10.2174/18752809787582561). [PubMed: [19275687](https://pubmed.ncbi.nlm.nih.gov/19275687/)].
- Ghanbari S, Yonessi M, Mohammadirad A, Gholami M, Baeeri M, Khorram-Khorshid HR, et al. Effects of IMOD and Angipars on mouse D-galactose-induced model of aging. *Daru*. 2012;**20**(1):68. doi: [10.1186/2008-2231-20-68](https://doi.org/10.1186/2008-2231-20-68). [PubMed: [23351487](https://pubmed.ncbi.nlm.nih.gov/23351487/)]. [PubMed Central: [PMC3555951](https://pubmed.ncbi.nlm.nih.gov/PMC3555951/)].
- Nickavar B, Mojab F, Javidnia K, Amoli MA. Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. *Z Naturforsch C*. 2003;**58**(9-10):629-31. doi: [10.1515/znc-2003-9-1004](https://doi.org/10.1515/znc-2003-9-1004). [PubMed: [14577620](https://pubmed.ncbi.nlm.nih.gov/14577620/)].
- Razavi BM, Hosseinzadeh H. A review of the effects of *Nigella sativa* L. and its constituent, thymoquinone, in metabolic syndrome. *J Endocrinol Invest*. 2014;**37**(11):1031-40. doi: [10.1007/s40618-014-0150-1](https://doi.org/10.1007/s40618-014-0150-1). [PubMed: [25125023](https://pubmed.ncbi.nlm.nih.gov/25125023/)].
- Forouzanfar F, Bazzaz BS, Hosseinzadeh H. Black cumin (*Nigella sativa*) and its constituent (thymoquinone): A review on antimicrobial effects. *Iran J Basic Med Sci*. 2014;**17**(12):929-38. [PubMed: [25859296](https://pubmed.ncbi.nlm.nih.gov/25859296/)]. [PubMed Central: [PMC4387228](https://pubmed.ncbi.nlm.nih.gov/PMC4387228/)].
- Hosseinzadeh H, Parvardeh S. Anticonvulsant effects of thymoquinone, the major constituent of *Nigella sativa* seeds, in mice. *Phytomedicine*. 2004;**11**(1):56-64. doi: [10.1078/0944-7113-00376](https://doi.org/10.1078/0944-7113-00376). [PubMed: [14971722](https://pubmed.ncbi.nlm.nih.gov/14971722/)].
- Amin B, Taheri MM, Hosseinzadeh H. Effects of intraperitoneal thymoquinone on chronic neuropathic pain in rats. *Planta Med*. 2014;**80**(15):1269-77. doi: [10.1055/s-0034-1383062](https://doi.org/10.1055/s-0034-1383062). [PubMed: [25272235](https://pubmed.ncbi.nlm.nih.gov/25272235/)].
- Hassan M, El-Dakhkhny M. Effect of some *Nigella sativa* constituents on chemical carcinogenesis in hamster cheek pouch. *Egypt Soci Pharmacol Exp Ther*. 1992;**11**:675-7.
- Houghton PJ, Zarka R, de las Heras B, Hoult JR. Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane lipid peroxidation. *Planta Med*. 1995;**61**(1):33-6. doi: [10.1055/s-2006-957994](https://doi.org/10.1055/s-2006-957994). [PubMed: [7700988](https://pubmed.ncbi.nlm.nih.gov/7700988/)].
- Marsik P, Kokoska L, Landa P, Nepovim A, Soudek P, Vanek T. In vitro inhibitory effects of thymol and quinones of *Nigella sativa* seeds on cyclooxygenase-1- and -2-catalyzed prostaglandin E2 biosyntheses. *Planta Med*. 2005;**71**(8):739-42. doi: [10.1055/s-2005-871288](https://doi.org/10.1055/s-2005-871288). [PubMed: [16142638](https://pubmed.ncbi.nlm.nih.gov/16142638/)].
- Hosseinzadeh H, Parvardeh S, Asl MN, Sadeghnia HR, Ziaee T. Effect of thymoquinone and *Nigella sativa* seeds oil on lipid peroxidation level during global cerebral ischemia-reperfusion injury in rat hippocampus. *Phytomedicine*. 2007;**14**(9):621-7. doi: [10.1016/j.phymed.2006.12.005](https://doi.org/10.1016/j.phymed.2006.12.005). [PubMed: [17291733](https://pubmed.ncbi.nlm.nih.gov/17291733/)].
- Mollazadeh H, Hosseinzadeh H. The protective effect of *Nigella sativa* against liver injury: A review. *Iran J Basic Med Sci*. 2014;**17**(12):958-66. [PubMed: [25859299](https://pubmed.ncbi.nlm.nih.gov/25859299/)]. [PubMed Central: [PMC4387231](https://pubmed.ncbi.nlm.nih.gov/PMC4387231/)].
- Bayrak O, Bavbek N, Karatas OF, Bayrak R, Catal F, Cimentepe E, et al. *Nigella sativa* protects against ischaemia/reperfusion injury in rat kidneys. *Nephrol Dial Transplant*. 2008;**23**(7):2206-12. doi: [10.1093/ndt/gfm953](https://doi.org/10.1093/ndt/gfm953). [PubMed: [18211980](https://pubmed.ncbi.nlm.nih.gov/18211980/)].
- Ismail N, Ismail M, Mazlan M, Latiff LA, Imam MU, Iqbal S, et al. Thymoquinone prevents beta-amyloid neurotoxicity in primary cultured cerebellar granule neurons. *Cell Mol Neurobiol*. 2013;**33**(8):1159-69. doi: [10.1007/s10571-013-9982-z](https://doi.org/10.1007/s10571-013-9982-z). [PubMed: [24101432](https://pubmed.ncbi.nlm.nih.gov/24101432/)].
- Abdollahi M, Momtaz S. A comprehensive review of biochemical and molecular evidences from animal and human studies on the role of oxidative stress in aging: An epiphenomenon or the cause. *Asia J Animal Vet Adv*. 2012;**7**(1):1-19. doi: [10.3923/ajava.2012.1.19](https://doi.org/10.3923/ajava.2012.1.19).
- Mahgoub AA. Thymoquinone protects against experimental colitis in rats. *Toxicol Lett*. 2003;**143**(2):133-43. doi: [10.1016/S0378-4274\(03\)00173-5](https://doi.org/10.1016/S0378-4274(03)00173-5). [PubMed: [12749817](https://pubmed.ncbi.nlm.nih.gov/12749817/)].
- Zhang ZF, Fan SH, Zheng YL, Lu J, Wu DM, Shan Q, et al. Purple sweet potato color attenuates oxidative stress and inflammatory response induced by d-galactose in mouse liver. *Food Chem Toxicol*. 2009;**47**(2):496-501. doi: [10.1016/j.fct.2008.12.005](https://doi.org/10.1016/j.fct.2008.12.005). [PubMed: [19114082](https://pubmed.ncbi.nlm.nih.gov/19114082/)].
- Yang YC, Lin HY, Su KY, Chen CH, Yu YL, Lin CC, et al. Rutin, a flavonoid that is a main component of *Saussurea involucrata*, attenuates the senescence effect in D-Galactose aging mouse model. *Evid Based Complement Alternat Med*. 2012;**2012**:980276. doi: [10.1155/2012/980276](https://doi.org/10.1155/2012/980276). [PubMed: [22952557](https://pubmed.ncbi.nlm.nih.gov/22952557/)]. [PubMed Central: [PMC3431096](https://pubmed.ncbi.nlm.nih.gov/PMC3431096/)].
- Jin SL, Yin YG. In vivo antioxidant activity of total flavonoids from *Indocalamus leaves* in aging mice caused by D-galactose. *Food Chem Toxicol*. 2012;**50**(10):3814-8. doi: [10.1016/j.fct.2012.07.046](https://doi.org/10.1016/j.fct.2012.07.046). [PubMed: [22874925](https://pubmed.ncbi.nlm.nih.gov/22874925/)].

20. Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem.* 1978;**86**(1):271-8. doi: [10.1016/0003-2697\(78\)90342-1](https://doi.org/10.1016/0003-2697(78)90342-1). [PubMed: [655387](https://pubmed.ncbi.nlm.nih.gov/655387/)].
21. Moron MS, Depierre JW, Mannervik B. Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver. *Biochim Biophys Acta.* 1979;**582**(1):67-78. doi: [10.1016/0304-4165\(79\)90289-7](https://doi.org/10.1016/0304-4165(79)90289-7). [PubMed: [760819](https://pubmed.ncbi.nlm.nih.gov/760819/)].
22. Fu H, Xu H, Chen H, Li Y, Li W, Zhu Q, et al. Inhibition of glycogen synthase kinase 3 ameliorates liver ischemia/reperfusion injury via an energy-dependent mitochondrial mechanism. *J Hepatol.* 2014;**61**(4):816-24. doi: [10.1016/j.jhep.2014.05.017](https://doi.org/10.1016/j.jhep.2014.05.017). [PubMed: [24862449](https://pubmed.ncbi.nlm.nih.gov/24862449/)].
23. Al-Johar D, Shinwari N, Arif J, Al-Sanea N, Jabbar AA, El-Sayed R, et al. Role of *Nigella sativa* and a number of its antioxidant constituents towards azoxymethane-induced genotoxic effects and colon cancer in rats. *Phytother Res.* 2008;**22**(10):1311-23. doi: [10.1002/ptr.2487](https://doi.org/10.1002/ptr.2487). [PubMed: [18570215](https://pubmed.ncbi.nlm.nih.gov/18570215/)].
24. Nagi MN, Alam K, Badary OA, al-Shabanah OA, al-Sawaf HA, al-Bekairi AM. Thymoquinone protects against carbon tetrachloride hepatotoxicity in mice via an antioxidant mechanism. *Biochem Mol Biol Int.* 1999;**47**(1):153-9. [PubMed: [10092955](https://pubmed.ncbi.nlm.nih.gov/10092955/)].
25. Zhou Y, Dong Y, Xu Q, He Y, Tian S, Zhu S, et al. Mussel oligopeptides ameliorate cognition deficit and attenuate brain senescence in D-galactose-induced aging mice. *Food Chem Toxicol.* 2013;**59**:412-20. doi: [10.1016/j.fct.2013.06.009](https://doi.org/10.1016/j.fct.2013.06.009). [PubMed: [23796539](https://pubmed.ncbi.nlm.nih.gov/23796539/)].
26. Kumar A, Prakash A, Dogra S. Centella asiatica attenuates D-galactose-induced cognitive impairment, oxidative and mitochondrial dysfunction in mice. *Int J Alzheimers Dis.* 2011;**2011**:347569. doi: [10.4061/2011/347569](https://doi.org/10.4061/2011/347569). [PubMed: [21629743](https://pubmed.ncbi.nlm.nih.gov/21629743/)]. [PubMed Central: [PMC3100561](https://pubmed.ncbi.nlm.nih.gov/PMC3100561/)].
27. Mehri S, Shahi M, Razavi BM, Hassani FV, Hosseinzadeh H. Neuroprotective effect of thymoquinone in acrylamide-induced neurotoxicity in Wistar rats. *Iran J Basic Med Sci.* 2014;**17**(12):1007-11. [PubMed: [25859305](https://pubmed.ncbi.nlm.nih.gov/25859305/)]. [PubMed Central: [PMC4387223](https://pubmed.ncbi.nlm.nih.gov/PMC4387223/)].
28. Brütüngaard H, Pedersen BK. Age-related inflammatory cytokines and disease. *Immunol Allergy Clin North Am.* 2003;**23**(1):15-39. doi: [10.1016/s0889-8561\(02\)00056-5](https://doi.org/10.1016/s0889-8561(02)00056-5).
29. Zhou SJ, Lu WH, Yuan D, Li H, Gu YQ, Wang CG, et al. [Changes of serum reproductive hormones with aging among healthy males in a community population of Hebei Province]. *Zhonghua Nan Ke Xue.* 2009;**15**(8):679-84. Chinese. [PubMed: [19852265](https://pubmed.ncbi.nlm.nih.gov/19852265/)].
30. Maggio M, Basaria S, Ceda GP, Ble A, Ling SM, Bandinelli S, et al. The relationship between testosterone and molecular markers of inflammation in older men. *J Endocrinol Invest.* 2005;**28**(11 Suppl Proceedings):116-9. [PubMed: [16760639](https://pubmed.ncbi.nlm.nih.gov/16760639/)].
31. Samir Bashandy AE. Effect of fixed oil of *nigella sativa* on male fertility in normal and hyperlipidemic rats. *Int J Pharmacol.* 2007;**3**(1):27-33. doi: [10.3923/ijp.2007.27.33](https://doi.org/10.3923/ijp.2007.27.33).