



Ozone Decreases Biomarkers of Inflammation (C-Reactive Protein and Erythrocyte Sedimentation Rate) and Improves Pain, Function and Quality of Life in Knee Osteoarthritis Patients: A Before-and-After Study and Review of the Literature

Marcos Edgar Fernandez-Cuadros,^{1,2,*} Olga Susana Perez-Moro,¹ Maria Jesus Albaladejo-Florin,¹ and Ruben Algarra-Lopez¹

¹Hospital Universitario Santa Cristina, Madrid, Spain

²Fundacion Hospital General Santisima Trinidad, Salamanca, Spain

*Corresponding author: Marcos Edgar Fernandez-Cuadros, Calle del Anzar, 44, piso Segundo, CP 28047, Madrid, Spain. Tel: +34-620314558, E-mail: marcosefc@hotmail.com

Received 2018 January 05; Revised 2018 February 19; Accepted 2018 February 21.

Abstract

Objectives: The aim of this study was to evaluate if ozone is capable of decreasing biomarkers of inflammation (CRP/ESR) and pain, and to improve function and quality of life in knee osteoarthritis (OA) patients.

Methods: A prospective quasi experimental (before-and-after study) research was performed to investigate the effect of Ozone therapy protocol. The intervention included 4 sessions (1 session/week) of an intra-articular infiltration of a medical mixture of Oxygen-Ozone (95% to 5%) at 20 ug/mL concentration. Biochemical evaluation (CRP and ESR), pain VAS, and WOMAC scales were evaluated before and after the treatment.

Results: Overall, 33 patients were evaluated. Mean age was 68.18 ± 8.43 years. Female patients accounted for 75.7% ($n = 25$) and male patients corresponded to 24.3% ($n = 8$). Female: male ratio was 3:1. The most frequent radiological KL grade was 2° ($n = 22$, 66.7%), followed by 3° ($n = 8$, 24.2%), and 4° ($n = 3$, 9.1%). Biomarkers of inflammation decreased significantly. C-reactive protein (CRP) diminished from 0.33 ± 0.32 mg/dL to 0.25 ± 0.23 mg/dL ($P = 0.0456$). Erythrocyte sedimentation rate (ESR) decreased from 15.06 ± 12.09 mm/h to 11.81 ± 8.32 mm/h ($P = 0.01$). Before treatment, pain measured by VAS was 7.33 ± 1.31 points and decreased to 2.84 ± 1.76 ($P < 0.0001$). The WOMAC-pain subscale was 14.84 ± 2.77 points and diminished to 5.96 ± 3.53 ($P < 0.0001$), WOMAC stiffness subscale was 3.06 ± 1.95 points and ameliorated to 1.15 ± 1.3 ($P < 0.0001$), WOMAC-function subscale was 41.15 ± 12.58 points and decreased to 22.3 ± 11.64 ($P < 0.0001$).

Conclusions: Ozone is capable of both decreasing pain and stiffness and improving function and quality of life in knee OA patients, yet decreases biomarkers of inflammation, such as CRP and ESR.

Keywords: Biochemical Markers, Ozone, Osteoarthritis, Inflammation, Acute-Phase Proteins, Blood Sedimentation, C-Reactive Proteins

1. Background

Osteoarthritis (OA) is the most prevalent joint disease. It is so common that almost 4 million people experience OA, and it is the cause of 50% total disability in Spain (1, 2). The economic impact is such that the direct cost of OA in Spain is 4738 million Euros per year. Therefore, OA becomes an important problem of public health (3). Moreover, OA produces pain, decreases function, and consumes a great amount of limited health resources. The emotional, social, and physical aspects are affected in patients, impairing their quality of life (QoL) (4).

Osteoarthritis influences cartilage at the involved joint; as a result, articular space narrows, and bone sclerosis, subchondral cysts, and osteophytes appear (4-7). Therefore, diagnosis is based on clinical and radiological signs and symptoms (8, 9). Knee OA is a multifactorial disease (4, 5). Risk factors include trauma, obesity, and recently, low-grade chronic inflammation (5-7, 10, 11).

The future challenge in the management of knee OA is to discover early tools for diagnosis, progression and monitoring of the disease, and to find effective therapeutic interventions. Biomarkers are among possible tools (12). Since OA is related to inflammation, inflamma-

tion biomarkers, such as C-reactive protein (CRP), Erythrocyte Sedimentation Rate (ESR), and pro-inflammatory cytokines (IL-1 β and IL-6) have been used to determine such relationship (8, 13-16).

In knee OA, the treatment goals in the short term are to ameliorate symptoms by symptomatic slow action drugs for OA (SYSADOA), and on the long term to diminish/revert articular damage and joint destruction by disease modifying drugs for OA (DMDOA) (3). The osteoarthritis research society international has suggested non-pharmacological, pharmacological, and surgical recommendations for the treatment/management of knee OA (2). Total knee arthroplasty is the last alternative in advanced knee OA patients, and although it is an invasive and expensive procedure with a 95% rate of survival at 10 years, it might have complications (4, 17, 18).

Several researchers worldwide and many years of clinical experience have indicated the fact that Ozone is capable to modulate inflammation (1, 19). In case of knee OA, several studies state that Ozone is capable of ameliorating pain and improving function and quality of life (SYSADOA effect) (2, 3, 5). One recent study by Fernandez-Cuadros et al., even stated that Ozone was capable of slowing/reverting knee OA progression in a 2-year follow-up study (DMDOA effect) (3). However, to the best of the author's knowledge, there is no study that has evaluated the effectiveness of ozone on biomarkers of inflammation, such as CRP and ESR on knee OA human subjects. There is only one report by Chang et al., who stated that Ozone could inhibit pro-inflammatory cytokines (IL1 and TNF- α) in an animal model (20).

The objective of this study was to evaluate if Ozone is capable of decreasing biomarkers of inflammation (CRP/ESR), ameliorating pain, and improving function and quality of life in knee OA patients.

2. Methods

A total of 33 patients were enrolled in a prospective quasi-experimental before-and-after study. Knee OA patients with a Kellgren-Lawrence (KL) grade of 2 or more, who attended the Santa Cristina's University hospital, with clinical symptoms (pain, stiffness, and loss of function), demanding conservative treatment, and in whom previous symptomatic treatment had failed, were included in the study. The study was conducted from January to November 2017, and it was authorized by the ethical committee of Santa Cristina's University hospital (Figure 1).

2.1. Inclusion Criteria

1, knee OA KL patients graded 2 or more; 2, pain greater than 3 on visual analogical scale (VAS), who had failed any

other conservative treatments (NAIDs, rehabilitation or physical therapy); 4, willingness to participate and not being a candidate for knee arthroplasty replacement.

2.2. Exclusion Criteria

1, allergy to Ozone (2, 3, 5); 2, incomplete biochemical analysis (CRP/ESR); 3, incomplete ozone therapy treatment protocol; and 4, absence from any of the questionnaires applied (VAS or WOMAC).

In the initial evaluation, the goals of treatment, procedure, indications and contraindications were explained, while initial biochemical evaluation (CRP and ESR analysis) was performed, informed consent was signed and outcome scales (VAS and WOMAC) were provided before the protocol (Figure 1).

The Ozone protocol consisted of 4 sessions (1 session/week) of an intra articular infiltration of a medical mixture of Oxygen-Ozone (95% to 5%) at a 20-ug/mL concentration. The skin area was cleaned with 1% chlorhexidine and anesthetized with ethyl chloride spray. The medical generator used was Ozonosan α -plus[®]. This equipment provided a 20-mL volume of 95% to 5% Oxygen-Ozone mixture that was introduced to a 20-mL silicone-coated syringe of 3 bodies. A Quincke needle of 27G, 4 cm was used to deliver Ozone to the joint.

Once the patient lied on the bed, the medical Ozone mixture was infiltrated on the knee on the superior and lateral side of the patella. If the infiltration is not performed in the joint, redness and pain appears after infiltration, however, these complaints disappear in a few minutes. The infiltration of Ozone treatment protocol on patient's knees was applied by the authors of the study.

After 4 sessions of ozone protocol, the final evaluation was accomplished, biomarker analysis, VAS and WOMAC Scales were applied, and adverse effects (if any) were registered. Serum levels of CRP and ESR were measured at baseline and at the end of the protocol treatment. The minimal measurable concentration of CRP is approximately 0.2 mg/dL and its coefficient of variation in expert technicians is 5.8% to 6.3% (21).

To perform the analysis of biochemical markers of inflammation, about 8 mL of venous blood was taken by arm venipuncture in a sterile vial. Five milliliters of blood was collected in a tube without anticoagulant, and serum was separated by centrifugation at 3500 rpm for 20 minutes. The level of CRP on serum was determined by the MULTIGENT CRP Vario assay (CRP Va), which is a latex immunoassay developed to measure accurately and reproducibly blood CRP levels in serum and plasma (CRP VARIO, Italy). The rest of the blood sample (3 mL) was collected in another sterile tube containing potassium EDTA anticoagulant for measurement of ESR by an automated sys-




Figure 1. Study design (n = 33). VAS, Visual Analogical Scale; WOMAC, Western Ontario and Mc Master University Index for Osteoarthritis; CRP, C-Reactive Protein; ESR, Erythrocyte Sedimentation Rate. The study ran from January to November 2017.

tem. The Ves-Matic Cube 30 (Diese, Italy) determines ESR, directly in closed tubes with EDTA blood samples using an infrared sensor to measure the level of opacity of the column of blood in the vertical position. The results obtained after 33 minutes of analysis corresponded to the Westergren method results after 60 minutes of sedimentation (Westergren method was the reference method).

The symptoms severity was measured by VAS and WOMAC scales. The VAS is a pain score graded from 0 to 10. A greater value is related to greater pain and vice versa (22). The WOMAC Index is a scale that evaluates pain, stiffness, and function, all in 24 questions. Each answer has 5 possible options: none, mild, moderately severe, and extreme. Pain includes 5 items (graded from 0 to 20), stiffness 2 items (graded from 0 to 8), and function 17 items (graded from 0 to 68) (22). Any change in the scores of the WOMAC index greater than 6% was considered as being clinically important. These changes represent, for WOMAC-pain subscale 1.2 points, for WOMAC-stiffness subscale 0.5 points, and for WOMAC-function subscale 4.1 points (23).

To radiologically classify knee OA, the KL scale was used. The KL grades are defined as follows: 0°, no radiological changes; 1°, doubtful osteophytes; 2°, osteophytes are present; 3°, joint space narrowing is observed; 4°, impingement, subchondral sclerosis, subchondral geodes, and marginal osteophytes are present.

Statistical analysis was performed by the use of SPSS® version 20.0. Frequencies and percentages were used to evaluate qualitative variables; while for the evaluation of quantitative variables, means and standard deviation were employed. The t-student test was the tool used for the evaluation of change before-and-after the treatment of quantitative variables. The X² test, if necessary, was employed to evaluate qualitative variables. The level of significance was 95% (P = 0.05).

3. Results

In this study, 33 patients were evaluated. Mean age of the patients was 68.18 ± 8.43 years. Female patients

Table 1. Principal Characteristics of Patients that Participated in the Study (N = 33)^{a, b}

Variables	Value
Age, y	68.18 ± 8.43
Female gender, n (%)	25 (75.7)
Male gender, n (%)	8 (24.3)
Female: Male ratio	3:1
OA KL 2°	22 (66.7)
OA KL 3°	8 (24.2)
OA KL 4°	3 (9.1)
CRP, mg/dL	0.33 ± 0.32
ESR, mm/h	15.06 ± 12.09
Pain VAS	7.33 ± 1.31
WOMAC pain score	14.84 ± 2.77
WOMAC stiffness score	3.06 ± 1.95
WOMAC function score	41.15 ± 12.58

Abbreviations: CRP, C-Reactive Protein; ESR, Erythrocyte Sedimentation Rate; KL, Kellgren-Lawrence; OA, Osteoarthritis; SD, Standard Deviation; VAS, Visual Analogical Scale; WOMAC, Western Ontario and Mc Master University Index for Osteoarthritis.

^aValues are expressed as No. (%) or mean ± SD.

^bP, T-student test.

accounted for 75.7% (n = 25), while male patients corresponded to 24.3% (n = 8), with a female:male ratio of 3:1 (Table 1).

The most frequent radiological KL grade was 2° (n = 22, 66.7%), followed by 3° grade (n = 8, 24.2%), and 4° grade (n = 3; 9.1%) (Table 1).

With respect to outcome variables, biomarkers of inflammation decreased significantly (P < 0.05) after Ozone therapy. The CRP diminished from 0.33 ± 0.32 mg/dL to 0.25 ± 0.23 mg/dL (P = 0.0456) (Table 2, Figure 2). The ESR decreased from 15.06 ± 12.09 mm/h to 11.81 ± 8.32 mm/h (P = 0.01) (Table 2, Figure 3).




Figure 2. Change of C-reactive protein (CRP) after Ozone therapy in knee osteoarthritis patients (n = 33)

Table 2. Effects of Ozone on biomarkers of Inflammation (CRP and ESR) and on Symptoms Severity (Measured by VAS and WOMAC Scales) in Knee OA Patients (n = 33)^a

Outcome Variables	Before	After	P Value ^b
CRP, mg/dL	0.33 ± 0.32	0.25 ± 0.23	0.0456
ESR, mm/h	15.06 ± 12.09	11.81 ± 8.32	0.0100
Pain VAS (0 - 10)	7.33 ± 1.31	2.84 ± 1.76	0.0000
WOMAC pain (0 - 20)	14.84 ± 2.77	5.69 ± 3.53	0.0000
WOMAC stiffness (0 - 8)	3.06 ± 1.95	1.15 ± 1.3	0.0000
WOMAC function (0 - 68)	41.15 ± 12.58	22.3 ± 11.64	0.0000

Abbreviations: CRP, C-Reactive Protein; ESR, Erythrocyte Sedimentation Rate; OA, Osteoarthritis; SD, Standard Deviation; VAS, Visual Analogical Scale; WOMAC, Western Ontario and Mc Master University Index for Osteoarthritis.

^aValues are expressed as mean ± SD.

^bT-student test.




Figure 3. Change of erythrocyte sedimentation rate (ESR) after Ozone therapy in knee osteoarthritis patients (n = 33)

Regarding symptom severity in knee OA, measured by VAS and WOMAC scales, Ozone therapy significantly improved each and every variable (P < 0.0001). Before treatment, pain measured by VAS was 7.33 ± 1.31 points and significantly decreased to 2.84 ± 1.76 points (P < 0.0001) (Table 2). Before the intervention, WOMAC-pain subscale was 14.84 ± 2.77 points, and diminished to 5.96 ± 3.53 points (P < 0.0001), WOMAC-stiffness subscale was 3.06 ± 1.95 points and decreased to 1.15 ± 1.3 points (P < 0.0001), and WOMAC-function subscale was 41.15 ± 12.58 points and improved to 22.3 ± 11.64 points (P < 0.0001) (Table 2).

4. Discussion

To the best of the author's knowledge, this is the first study that stated that Ozone decreases biomarkers of inflammation, namely CRP and ESR in human knee OA patients, and the diminishing comes in line with a clinical improvement on symptoms severity, such as pain, stiffness, function, and QoL.

In the current study, and because of economical limitations, the researchers intended to evaluate the effectiveness of Ozone on knee OA patients by the use of acute phase proteins/reactants of inflammation, a hallmark of knee OA (1, 5, 24-33). These biomarkers are cheap, easy to perform, and widely used in clinical practice. The objective of this study was to demonstrate that CRP/ESR biomarkers of inflammation decrease after Ozone treatment, since it was hypothesized that Ozone improves pain, function, stiffness and QoL, and many studies suggest that ozone is capable of ameliorating inflammation (5). The rationale to evaluate CRP/ESR is that inflammation biomarkers may predict outcomes in OA. Elevated hs-CRP predicts loss of cartilage in knee OA and poor results after total knee arthroplasty.

Biomarkers of inflammation used in the study were CRP and ESR. These biomarkers are recognized as acute phase reactants or proteins. They indicate an inflammatory state, as in the case of rheumatic diseases. Erythrocyte Sedimentation Rate is cheap and commonly used in clinical practice. It mainly reflects fibrinogen concentration. Furthermore, CRP shows no variation with age, neither with erythrocytes morphology, nor with other protein variations. During inflammation, CRP concentration increases by 1000 folds; on the contrary, ESR increases by 2 to 5 folds. In an inflammatory process, such as orthopedic surgery, CRP reaches its maximum level at 24 to 48 hours and normalizes by the next week. On the contrary, ESR value is elevated after a week, and normalizes in the next 6 weeks (13).

The current findings and several other facts suggest that the development of OA is usually accompanied by inflammation (16). Epidemiological studies suggest that the severity and progression of tibiofemoral cartilage loss are more frequent and severe in patients with inflamed synovial fluid (34). Higher levels of IL-1 β and TNF- α , which are mediators of inflammation, are found in the progression of the disease (16). Several studies have observed that increased CRP levels are related to prevalence and progression of the knee or hip OA (16). With respect to OA, elevated CRP levels are related to synovial fluid IL-6 levels and to synovial infiltration, as well as with symptoms of pain and stiffness, radiographic grading, and disease progression (35).

Many researchers worldwide have suggested that not only IL-6, yet CRP and ESR are closely related to radiographic knee OA, symptoms severity, and the progression of the disease (8, 36-44). These biomarkers suggest that low-grade inflammation might be a direct pathway in structural and symptomatic changes on knee OA (21).

All the presented data suggest that inflammatory biomarkers, such as CRP and ESR, might be a key factor in knee OA pathogenesis; they could serve as OA predic-

tors and as outcome variables. Their change may indicate the effectiveness of the current intervention (Ozone treatment). With this in mind, it could be understood why in the present study CRP and ESR were slightly elevated; and how Ozone was capable of diminishing such biochemical levels, together with symptom relief. This observation also comes in line with previous studies (Fernandez-Cuadros et al), in which important and significant amelioration of pain, rigidity; and improvement on QoL was observed after Ozone treatment. Indeed, for the first time, the findings correlate clinical improvement with biochemical improvement in knee OA patients. The current study is in accordance with that of Zhu et al., who stated that hs-CRP was significantly associated with WOMAC knee pain and its changes (21).

From the previous discussion, it can be inferred that the decrease of CRP and ESR as markers of inflammation, reflect the effectiveness of Ozone treatment on knee OA management. Therefore, in the current study it was observed that Ozone ameliorates pain and stiffness, improves pain and function, and decreases CRP and ESR values. This study also showed a direct relationship between inflammation and symptom severity in knee OA. Before treatment, patients presented a high score on VAS and WOMAC scales, and high values of serum CRP and ESR. After the intervention (Ozone treatment), lower scores on VAS and WOMAC scales were observed, which correspond to symptoms alleviation, and were related to lower CRP and ESR values.

An important limitation of the study was the absence of a control group. The small sample size ($n = 33$) limited the establishment of a control group. Indeed, because of ethical considerations, the researchers pursued to treat all patients with the treatment protocol, based on decades of clinical experience in the management of Ozone for pain and inflammation. This quasi-experimental before-and-after study intended to add clinical-based evidence and solve the lack of control group. That is the main reason why this kind of study was designed. In the before-and-after evaluation, a change after the intervention was expected to be the direct effect of Ozone on the treatment group.

Another important limitation subsides in the fact that the specificity of biomarkers of inflammation on knee OA was limited, because they are compromised in several inflammatory disorders. Despite much active research on biomarkers in OA, no single biomarker has been found to be the gold standard or sufficiently well validated and recognized as a valuable tool for diagnosis, prognosis, investigation and/or evaluation of effectiveness on knee OA (8).

Economic constraints limited the use/evaluation of CRP/ESR as general markers of inflammation. Despite the sample size, biomarker specificity of CRP and ESR, and the

economic limitations of this study, such limitations do not affect the observations obtained from this study. The reduction of inflammation biomarkers means that inflammation might constitute an important factor in the progression of knee OA, although further research on this issue must be accomplished. Future studies should be performed on the effectiveness of Ozone to reduce specific inflammatory cytokines in serum or synovial fluid, such as Interleukin (IL)-1, IL-6, Tumor Necrosis Factor (TNF)- α , and Interferon (IFN)- γ .

4.1. Conclusion

In the light of the current results, it could be concluded that Ozone is capable not only of decreasing pain and stiffness and improving function and quality of life in knee OA patients, yet it could also decrease biomarkers of inflammation such as CRP and ESR.

Acknowledgments

To Aureliana Gomez Sierra for the collection of data and preparation of material for intra articular infiltration, to María del Carmen de la Riva Molina and to Saturnino Diaz-Trujillo for the article search.

References

- Fernandez-Cuadros ME, Perez-Moro OS, Albaladejo-Florin MJ. *Ozone Fundamentals and effectiveness on knee pain: Chondromalacia and knee Osteoarthritis*. Germany: Lambert Academic Publishing; 2016.
- Fernández Cuadros ME, Pérez Moro OS, Albaladejo Florin MJ, Mirón Canelo JA. Ozone improves pain, function and quality of life in patients with knee osteoarthritis: A prospective quasi-experimental before-after study. *Middle East J Rehabil Health*. 2016;**4**(1). doi: [10.17795/mejrh-41821](#).
- Fernandez-Cuadros ME, Perez-Moro OS, Albaladejo-Florin MJ. Knee osteoarthritis: Chondroprotector action and symptomatic effect of ozone on pain, function, quality of life, minimal joint space and knee arthroplasty delay. *Middle East J Rehabil Health*. 2017;**4**(1).
- Fernández-Cuadros ME. *Análisis de la calidad de vida en pacientes con prótesis total de rodilla*. Spain: Universidad de Salamanca; 2013.
- Fernandez-Cuadros ME, Perez-Moro OS, Miron-Canelo JA. Could Ozone be used as a feasible future treatment in osteoarthritis of the knee? *Divers Equal Health Care*. 2016;**13**(3):232-9.
- Mobasher A, Bay-Jensen AC, van Spil WE, Larkin J, Levesque MC. Osteoarthritis Year in Review 2016: biomarkers (biochemical markers). *Osteoarthritis Cartilage*. 2017;**25**(2):199-208. doi: [10.1016/j.joca.2016.12.016](#). [PubMed: [28099838](#)].
- Vaillant JD, Fraga A, Diaz MT, Mallok A, Viebahn-Hansler R, Fahmy Z, et al. Ozone oxidative postconditioning ameliorates joint damage and decreases pro-inflammatory cytokine levels and oxidative stress in PG/PS-induced arthritis in rats. *Eur J Pharmacol*. 2013;**714**(1-3):318-24. doi: [10.1016/j.ejphar.2013.07.034](#). [PubMed: [23911887](#)].
- Lotz M, Martel-Pelletier J, Christiansen C, Brandi ML, Bruyere O, Chapurlat R, et al. Value of biomarkers in osteoarthritis: current status and perspectives. *Ann Rheum Dis*. 2013;**72**(11):1756-63. doi: [10.1136/annrheumdis-2013-203726](#). [PubMed: [23897772](#)].
- Rubén Daniel Arellano PV, Jesús Rafael AA, Morán-Martínez J, García-Marín A, Guzmán D, Lizette Sarai AM. Biochemical Markers in Osteoarthritis. *Int J Bone Rheumatol Res*. 2015;**2**(2):11-7. doi: [10.19070/2470-4520-150003](#).
- Weinstein AM, Rome BN, Reichmann WM, Collins JE, Burbine SA, Thornhill TS, et al. Estimating the burden of total knee replacement in the United States. *J Bone Joint Surg Am*. 2013;**95**(5):385-92. doi: [10.2106/JBJS.L.00206](#). [PubMed: [23344005](#)].
- Alberto Alexandre EB, Andrea Alexandre EI. Disc herniation and knee arthritis as chronic oxidative stress diseases: The therapeutic role of Oxygen Ozone therapy. *J Arthritis*. 2015;**4**(3). doi: [10.4172/2167-7921.1000161](#).
- Möller I, Gharbi M, Martínez H, Herrero M, Vergés J, Henrotin Y. Effect of chondroitin sulfate on soluble biomarkers of osteoarthritis: How to analyze and interpret the results from an open-label trial in unilateral knee osteoarthritis patients. *Osteoarthr Cartil*. 2016;**24**:77. doi: [10.1016/j.joca.2016.01.165](#).
- García FJB. *Manual SER de las enfermedades reumáticas*. Médica Panamericana; 2004.
- Saxne T, Lindell M, Mansson B, Petersson IF, Heinegard D. Inflammation is a feature of the disease process in early knee joint osteoarthritis. *Rheumatology (Oxford)*. 2003;**42**(7):903-4. doi: [10.1093/rheumatology/keg214](#). [PubMed: [12826709](#)].
- Sowers M, Jannausch M, Stein E, Jamadar D, Hochberg M, Lachance L. C-reactive protein as a biomarker of emergent osteoarthritis. *Osteoarthritis Cartilage*. 2002;**10**(8):595-601. doi: [10.1053/joca.2002.0800](#). [PubMed: [12479380](#)].
- Malathi R. Raised serum IL 6 and CRP in radiographic knee osteoarthritis in Eastern India. *J Med Sci Clin Res*. 2017;**5**(5):21687-92. doi: [10.18535/jmscr/v5i5.73](#).
- Fernandez-Cuadros ME, Pérez-Moro OS, Miron-Canelo JA. Knee osteoarthritis: Impact on quality of life and effectiveness of total knee arthroplasty. *Divers Equal Health Care*. 2016;**13**(4):278-83.
- Fernandez-Cuadros ME, Perez-Moro OS, Miron-Canelo JA. Age and comorbidities affect quality of life in patients with osteoarthritis and knee replacement. *Middle East J Rehabil Health*. 2016;**3**(4). doi: [10.17795/mejrh-40247](#).
- Mawsouf MN, El-Sawalhi MM, Darwish HA, Shaheen AA, Martínez-Sánchez G, Re L. Effect of ozone therapy on redox status in experimentally induced arthritis. *Revista Española de Ozonoterapia*. 2012;**1**(1):32-4.
- Chang JD, Lu HS, Chang YF, Wang D. Ameliorative effect of ozone on cytokine production in mice injected with human rheumatoid arthritis synovial fibroblast cells. *Rheumatol Int*. 2005;**26**(2):142-51. doi: [10.1007/s00296-004-0526-1](#). [PubMed: [15570426](#)].
- Zhu Z, Jin X, Wang B, Wluka A, Antony B, Laslett LL, et al. Cross-Sectional and Longitudinal Associations Between Serum Levels of High-Sensitivity C-Reactive Protein, Knee Bone Marrow Lesions, and Knee Pain in Patients With Knee Osteoarthritis. *Arthritis Care Res (Hoboken)*. 2016;**68**(10):1471-7. doi: [10.1002/acr.22834](#). [PubMed: [26748954](#)].
- Perez-Moro O, Albaladejo-Florin M, Entrambasaguas-Esteba B, Fernandez-Cuadros M. Effectiveness of PRP on pain, function and quality of life in chondromalacia and patellofemoral pain syndrome: A pretest-posttest analysis. *Nov Tech Arthritis Bone Res*. 2017;**1**(1).
- Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*. 1988;**15**(12):1833-40. [PubMed: [3068365](#)].
- Kraus VB, Blanco FJ, Englund M, Karsdal MA, Lohmander LS. Call for standardized definitions of osteoarthritis and risk stratification for clinical trials and clinical use. *Osteoarthritis Cartilage*. 2015;**23**(8):1233-41. doi: [10.1016/j.joca.2015.03.036](#). [PubMed: [25865392](#)].
- Rainbow R, Ren W, Zeng L. Inflammation and Joint Tissue Interactions in OA: Implications for Potential Therapeutic Approaches. *Arthritis*. 2012;**2012**:741582. doi: [10.1155/2012/741582](#). [PubMed: [22745906](#)].

26. Scanzello CR, Goldring SR. The role of synovitis in osteoarthritis pathogenesis. *Bone*. 2012;**51**(2):249–57. doi: [10.1016/j.bone.2012.02.012](https://doi.org/10.1016/j.bone.2012.02.012). [PubMed: [22387238](https://pubmed.ncbi.nlm.nih.gov/22387238/)].
27. Berenbaum F. Osteoarthritis as an inflammatory disease (osteoarthritis is not osteoarthritis!). *Osteoarthritis Cartilage*. 2013;**21**(1):16–21. doi: [10.1016/j.joca.2012.11.012](https://doi.org/10.1016/j.joca.2012.11.012). [PubMed: [23194896](https://pubmed.ncbi.nlm.nih.gov/23194896/)].
28. Nguyen LT, Sharma AR, Chakraborty C, Saibaba B, Ahn ME, Lee SS. Review of Prospects of Biological Fluid Biomarkers in Osteoarthritis. *Int J Mol Sci*. 2017;**18**(3). doi: [10.3390/ijms18030601](https://doi.org/10.3390/ijms18030601). [PubMed: [28287489](https://pubmed.ncbi.nlm.nih.gov/28287489/)].
29. Griffin TM, Huebner JL, Kraus VB, Yan Z, Guilak F. Induction of osteoarthritis and metabolic inflammation by a very high-fat diet in mice: effects of short-term exercise. *Arthritis Rheum*. 2012;**64**(2):443–53. doi: [10.1002/art.33332](https://doi.org/10.1002/art.33332). [PubMed: [21953366](https://pubmed.ncbi.nlm.nih.gov/21953366/)].
30. Muschter D, Gottl C, Vogel M, Grifka J, Straub RH, Grassel S. Reactivity of rat bone marrow-derived macrophages to neurotransmitter stimulation in the context of collagen II-induced arthritis. *Arthritis Res Ther*. 2015;**17**:169. doi: [10.1186/s13075-015-0684-4](https://doi.org/10.1186/s13075-015-0684-4). [PubMed: [26104678](https://pubmed.ncbi.nlm.nih.gov/26104678/)].
31. Benito MJ, Veale DJ, FitzGerald O, van den Berg WB, Bresnihan B. Synovial tissue inflammation in early and late osteoarthritis. *Ann Rheum Dis*. 2005;**64**(9):1263–7. doi: [10.1136/ard.2004.025270](https://doi.org/10.1136/ard.2004.025270). [PubMed: [15731292](https://pubmed.ncbi.nlm.nih.gov/15731292/)].
32. Sun AR, Panchal SK, Friis T, Sekar S, Crawford R, Brown L, et al. Obesity-associated metabolic syndrome spontaneously induces infiltration of pro-inflammatory macrophage in synovium and promotes osteoarthritis. *PLoS One*. 2017;**12**(8). e0183693. doi: [10.1371/journal.pone.0183693](https://doi.org/10.1371/journal.pone.0183693). [PubMed: [28859108](https://pubmed.ncbi.nlm.nih.gov/28859108/)].
33. Rousseau J, Garnero P. Biological markers in osteoarthritis. *Bone*. 2012;**51**(2):265–77. doi: [10.1016/j.bone.2012.04.001](https://doi.org/10.1016/j.bone.2012.04.001). [PubMed: [22538364](https://pubmed.ncbi.nlm.nih.gov/22538364/)].
34. Ayril X, Pickering EH, Woodworth TG, Mackillop N, Dougados M. Synovitis: a potential predictive factor of structural progression of medial tibiofemoral knee osteoarthritis – results of a 1 year longitudinal arthroscopic study in 422 patients. *Osteoarthritis Cartilage*. 2005;**13**(5):361–7. doi: [10.1016/j.joca.2005.01.005](https://doi.org/10.1016/j.joca.2005.01.005). [PubMed: [15882559](https://pubmed.ncbi.nlm.nih.gov/15882559/)].
35. Smith JW, Martins TB, Gopez E, Johnson T, Hill HR, Rosenberg TD. Significance of C-reactive protein in osteoarthritis and total knee arthroplasty outcomes. *Ther Adv Musculoskelet Dis*. 2012;**4**(5):315–25. doi: [10.1177/1759720X12455959](https://doi.org/10.1177/1759720X12455959). [PubMed: [23024709](https://pubmed.ncbi.nlm.nih.gov/23024709/)].
36. Spector TD, Hart DJ, Nandra D, Doyle DV, Mackillop N, Gallimore JR, et al. Low-level increases in serum C-reactive protein are present in early osteoarthritis of the knee and predict progressive disease. *Arthritis Rheum*. 1997;**40**(4):723–7. doi: [10.1002/1529-0131\(199704\)40:4<723::AID-ART18>3.0.CO;2-L](https://doi.org/10.1002/1529-0131(199704)40:4<723::AID-ART18>3.0.CO;2-L). [PubMed: [9125256](https://pubmed.ncbi.nlm.nih.gov/9125256/)].
37. Pelletier JP, Raynaud JP, Caron J, Mineau F, Abram F, Dorais M, et al. Decrease in serum level of matrix metalloproteinases is predictive of the disease-modifying effect of osteoarthritis drugs assessed by quantitative MRI in patients with knee osteoarthritis. *Ann Rheum Dis*. 2010;**69**(12):2095–101. doi: [10.1136/ard.2009.122002](https://doi.org/10.1136/ard.2009.122002). [PubMed: [20570834](https://pubmed.ncbi.nlm.nih.gov/20570834/)].
38. Pearle AD, Scanzello CR, George S, Mandl LA, DiCarlo EF, Peterson M, et al. Elevated high-sensitivity C-reactive protein levels are associated with local inflammatory findings in patients with osteoarthritis. *Osteoarthritis Cartilage*. 2007;**15**(5):516–23. doi: [10.1016/j.joca.2006.10.010](https://doi.org/10.1016/j.joca.2006.10.010). [PubMed: [17157039](https://pubmed.ncbi.nlm.nih.gov/17157039/)].
39. Jin X, Beguerie JR, Zhang W, Blizzard L, Otahal P, Jones G, et al. Circulating C reactive protein in osteoarthritis: a systematic review and meta-analysis. *Ann Rheum Dis*. 2015;**74**(4):703–10. doi: [10.1136/annrheumdis-2013-204494](https://doi.org/10.1136/annrheumdis-2013-204494). [PubMed: [24363360](https://pubmed.ncbi.nlm.nih.gov/24363360/)].
40. Saberi Hosnijeh F, Siebuhr AS, Uitterlinden AG, Oei EH, Hofman A, Karsdal MA, et al. Association between biomarkers of tissue inflammation and progression of osteoarthritis: evidence from the Rotterdam study cohort. *Arthritis Res Ther*. 2016;**18**:81. doi: [10.1186/s13075-016-0976-3](https://doi.org/10.1186/s13075-016-0976-3). [PubMed: [27039382](https://pubmed.ncbi.nlm.nih.gov/27039382/)].
41. He Y, Siebuhr AS, Brandt-Hansen NU, Wang J, Su D, Zheng Q, et al. Type X collagen levels are elevated in serum from human osteoarthritis patients and associated with biomarkers of cartilage degradation and inflammation. *BMC Musculoskelet Disord*. 2014;**15**:309. doi: [10.1186/1471-2474-15-309](https://doi.org/10.1186/1471-2474-15-309). [PubMed: [25245039](https://pubmed.ncbi.nlm.nih.gov/25245039/)].
42. Sanchez-Ramirez DC, van der Leeden M, van der Esch M, Gerritsen M, Roorda LD, Verschueren S, et al. Association of serum C-reactive protein and erythrocyte sedimentation rate with muscle strength in patients with knee osteoarthritis. *Rheumatology (Oxford)*. 2013;**52**(4):727–32. doi: [10.1093/rheumatology/kes366](https://doi.org/10.1093/rheumatology/kes366). [PubMed: [23275388](https://pubmed.ncbi.nlm.nih.gov/23275388/)].
43. Stannus OP, Jones G, Blizzard L, Cicuttini FM, Ding C. Associations between serum levels of inflammatory markers and change in knee pain over 5 years in older adults: a prospective cohort study. *Ann Rheum Dis*. 2013;**72**(4):535–40. doi: [10.1136/annrheumdis-2011-201047](https://doi.org/10.1136/annrheumdis-2011-201047). [PubMed: [22580582](https://pubmed.ncbi.nlm.nih.gov/22580582/)].
44. Arendt-Nielsen L, Eskehave TN, Egsgaard LL, Petersen KK, Graven-Nielsen T, Hoeck HC, et al. Association between experimental pain biomarkers and serologic markers in patients with different degrees of painful knee osteoarthritis. *Arthritis Rheumatol*. 2014;**66**(12):3317–26. doi: [10.1002/art.38856](https://doi.org/10.1002/art.38856). [PubMed: [25168637](https://pubmed.ncbi.nlm.nih.gov/25168637/)].