



Repeated Training Creates Spatial Memory in an Adult Male Rat Model of Testosterone-Induced Spatial Learning Impairment

Azadeh Gholaminejad¹, Nasser Naghdi^{1,2}, Hamid Gholamipour-Badie² and Mohammad Nasehi^{1,3,*}

¹Department of Cognitive Neuroscience, Institute for Cognitive Science Studies, Tehran, Iran

²Department of Physiology and Pharmacology, Pasteur Institute of Iran, Tehran, Iran

³Cognitive and Neuroscience Research Center, Amir-Almomenin Hospital, Islamic Azad University, Tehran, Iran

*Corresponding author: Cognitive and Neuroscience Research Center, Islamic Azad University, Amir-Almomenin Hospital, Tehran, Iran. Tel: +98-9128224672, Fax: +98-2199881117, Email: nasehi@iricss.org

Received 2019 January 06; Revised 2019 February 10; Accepted 2019 February 13.

Abstract

Background: Memories are primarily defined as the fragmentary or partial reconstruction of what were actually experienced at the time of acquisition. Spatial learning is assessed through repeated trials and reference memory is assessed by the measurement of latency in finding a concealed platform preference for the platform area when the platform is not present. Our daily experience, as well as learning experiments performed in animal studies, has enabled us to know that the formation of long-lasting memory needs repeated practice.

Objectives: Our study aimed to investigate the effect of repeated training on spatial learning impairment, which was induced by the administration of testosterone in young adult rats.

Methods: Cannula were bilaterally implanted into the Cornu Ammon (CA1) region of the hippocampus while testosterone (Testosterone Enantate, Aburaihan Pharmaceutical Company, Tehran) was daily microinjected for 3 minutes in each side. In this study, twenty-four male adult rats were divided into three groups as follows: the control group that received no treatment, the sham group that received DMSO as a drug solvent, and the treatment group that received testosterone at a dose of 80 µg/0.5 µL DMSO/each side injected into the CA1 before each session.

Results: The results showed that the bilateral administration of testosterone into the CA1 region significantly increased the escape latency and the distance traveled by rats compared with the control and sham groups in the acquisition test. However, in the probe test (retrieval) there was no difference between the treatment group and other groups considering the escape latency and traveled distance.

Conclusions: It seems that intra CA1 microinjection of testosterone causes the impairment in spatial learning. Repeated training enhances spatial learning in the Morris water maze (MWM) task, which leads to repeated long-term potentiation (LTP), spinogenesis, increased spine density, and spontaneous generation of new spines, resulting in the improvement of spatial memory in retrieval test.

Keywords: CA1, Learning Impairment, Morris Water Maze, Repeated Training, Testosterone

1. Background

The conception of memory along with its entity has not been fully elucidated as the molecular mechanisms underlying the storage of memory, and the physical nature of its trace remained unclear. While numerous efforts have been made to unravel the process of memory storage and function, the only reliable method to detect such phenomena is the monitoring of behavioral changes in animal studies to get a new insight, yet indirect, on the functionality of the memory (1). The Morris water maze (MWM) task is a behavioral test used for the determination of spatial learning in rodents. Spatial learning is assessed through repeated trials and reference memory is monitored via preference

for the platform area once the platform is missing (2). The obtained information about our daily experience and the establishment of learning experiments in animal studies have revealed that the formation of the long-lasting memory needs repeated practice (3). Spatial memory formation in the MWM task mainly depends upon the hippocampus function in animals. A growing body of evidence indicates that the hippocampus is vital for the acquisition and retrieval of spatial information (4). Studies have demonstrated that the performance in MWM task is tightly associated with hippocampal synaptic plasticity, as measured via long-term potentiation (LTP), and the function of NMDA. These properties have made the MWM a vigorous and re-

liable examination (5). Myriad investigations have highlighted that the NMDA receptors, which are present in the brain, are involved in many essential biological functions such as neurotoxicity, neuronal plasticity, as well as memory and learning (6). It should be noted that the memory is constituted of multiple stages, including acquisition, consolidation, and retrieval. However, the separation and characterization of each step are experimentally tricky (7). One of the outcomes of the consolidation paradigm is that the experimental mediation could be implemented in two or three different moments during memory formation and recall. Because the formation procedure is not merely an instant process, three formation phases could be determined, namely acquisition which is known as learning, and consolidation, the unstable phase in which the memory stabilizes, and memory retrieval, the stage of bringing back of the learned task (8). Experimental research has reported that the probe trial performance is able to evaluate the spatial memory of laboratory animals (9). One of the conceivable facets of the memory function is the acquisition stage as some molecules have been attributed to this stage of the memory. One of the popular aspects of the acquisition stage of the memory is the availability of the cellular models of learning, involving various forms of synaptic plasticity such as LTP (10). A number of studies implicated that LTP is supposed to be correlated with learning and memory and, more generally stated, enduring experience-dependent improvement of synaptic transmission (5) Hence, the primary target for excitatory synaptic contact dendritic spines, which are located on the pyramidal neurons, hippocampus, and neocortex. Basically, dendritic spines provide a site of synaptic contact by which the neuronal cells could receive input from another neuron (11). It has been shown that dendritic spine density is increased on pyramidal cells in the Cornu Ammon (CA1) region following of two different spatial memory tasks, the MWM and object placement suggesting that they are morphological substrates for the memory function (12). On the other hand, studies indicate that the effects of androgens on the density of spine synapses located on pyramidal neurons in the CA1 area of the hippocampus in male rats. In the CA1 region, the androgen receptors were mainly located on pyramidal neurons, suggesting a potential target for the testosterone action. Gonadectomy had no considerable effects on the number of CA1 pyramidal cells but decreased the CA1 spine synapse density by nearly 50% in comparison with sham and control groups. The treatment of castrated rats with testosterone propionate the density of spine synapse was increased to levels in comparison to the healthy male rats. A similar increase was observed in synapse density in castrated rats after treatment with dihydrotestosterone (DHT) (13).

2. Methods

2.1. Animals

Male Wistar rats weighing 200 - 250 g were purchased from the Pasteur Institute of Karaj, Tehran, Iran. The male rats were housed before the surgical procedures at $23 \pm 2^\circ\text{C}$ and the humidity ($50 \pm 5\%$) with a 12-h light:12-h dark cycle.

2.2. Surgical Procedures

Rats were anesthetized with intraperitoneal administration of ketamine and xylazine, Alfasan Company, Netherlands (3 mg/100 g and 1mg/100 g of body weight, respectively) and placed in a stereotaxic instrument. Then, two guide cannula were bilaterally implanted in the CA1 region of the hippocampus at coordinates: AP= -3.8mm, ML= ± 2.2 mm from Bregma and D/V= -2.7mm from dura (4).

2.3. Microinjection Procedure

The intracerebral injection was administered by means of guide cannula using injection needles connected to 10- μL Hamilton microsyringes by polyethylene tubing. The administration of DMSO (0 μg testosterone, 0.5 μL DMSO/side; 30 min before training) and testosterone (80 μg , 0.5 μL DMSO/side; 30 min before training) was performed (pre-training) over 3 min.

2.4. Behavioral Assessment Apparatus

The MWM task consisted of a dark round pool that was filled with water. A transparent platform was placed 1 cm below the surface of the water at the center of the arbitrarily designed north-east, south-east, south-west or north-west orthogonal quadrant. The pool was located in a specific room specified for behavioral experiments with a constant environment and visual cues on curtains around the pool. The animals were placed in water and positioned to face the wall of the pool, and allowed to find the hidden platform underwater (4). A video camera connected to a tracking device (Ethovision XT, version 7) was placed above the pool to monitor the analysis of each trial and measure the traveled distance, escape latency, and swimming speed.

2.5. Behavioral Assessment

In the hidden platform test, the animals were submitted to a daily session of four trials for four consecutive days. During each trial, ninety seconds were specified for each rat to find the platform positioned at the center of the target quadrant. Rats were allowed to remain there for twenty seconds after each trial; then the next trial started and new starting points changed in each trial by a stochastic process. On the fifth day of the experiment, the platform was removed in order to perform the retrieval learning and

memory test and the rats were released from the opposite location and allowed to swim freely for sixty seconds. In the visible platform test, the platform was elevated above the surface of the water in order to be visible and then it was placed in different positions. The animals were requested to swim in water at four different directions of the tank in four trials. The obtained results were evaluated and extrapolated by the visio-motor coordination and motivational system.

2.6. Statistical Analysis

The statistical evaluations were carried out using the SPSS software version 23.0. The data were analyzed using one-way analysis of variance (One-way ANOVA) and repeated measures (RM) two-way analysis of variance (Two-way ANOVA) followed by Tukey's post hoc test. The difference between the groups was statistically significant when the p-value was less than 0.05.

2.7. Study Design

In this study, twenty-four male rats were divided into three groups. The first group was designated as the control group that received no treatment but was trained every day. The second group was coined as the sham group that received DMSO as a drug solvent. The third group (treatment group) received testosterone at a dose of 80 $\mu\text{g}/0.5 \mu\text{L}$ DMSO/each side, dissolved in 0.5 μL vehicle (DMSO). Both of the sham and treatment groups were daily microinjected 30 min before the hidden platform test. The purpose of the experiment was to determine the impact of bilateral injection of testosterone into the CA1 region on the spatial learning and memory outcomes.

3. Results

3.1. Microinjection of Testosterone Into the CA1 Region Induced Spatial Learning Impairment

Figure 1 shows the results obtained from the injection of testosterone. The results indicated a significant effect of treatment [$F(2,80) = 23.04$; $P < 0.0001$] on the morphological changes in the CA1 region of the brain. There was also a significant increase in the escape latency in the first ($P < 0.001$), second, and third ($P < 0.01$) training days in the treatment group compared with the control group. Also, there was a significant increase in the escape latency in the first ($P < 0.05$) and second ($P < 0.01$) days in the treatment group when compared with the sham group (Figure 1A). Also, the findings showed a significant impact of testosterone administration [$F(2, 80) = 19.55$; $P < 0.001$] on travel distance parameters. The statistical analysis demonstrated a significant increment in the traveled distance in the first ($P < 0.001$), second, and third ($P < 0.01$) training days in the

testosterone-treated group in comparison to the control group. Similarly, there was a significant increase in traveled distance in the first ($P < 0.01$) and second ($P < 0.05$) training days in the treatment group compared with the sham group (Figure 1B). There was no significant difference between the testosterone-treated group and other groups when the swimming speed was compared in all training days (Figure 1C).

Figure 2 shows the comparative effects of testosterone injection in all experimental groups. A significant increase was found in the escape latency [$F(2,20) = 13.942$; $P = 0.000$] and traveled distance [$F(2,20) = 10.247$; $P = 0.001$] among the experimental groups (Figure 2A and B). No significant difference was observed among the studied groups considering the swimming speed (Figure 2C).

According to Figure 3, there was no significant difference between the treatment group and other groups regarding the time spent in the target quadrant (Figure 3A). Accordingly, there was no significant difference among all groups in terms of the traveled distance to the platform (Figure 3B). Of note, there was no significant difference among all groups of the study considering the escape latency (Figure 3C) or velocity (Figure 3D) in the visible platform test.

4. Discussion

In the present study, we found that microinjection of testosterone (80 $\mu\text{g}/0.5 \mu\text{L}$ DMSO/each side) into the CA1 region resulted in impairment in spatial learning but not in spatial memory and this may contribute to the memory formation. The behavioral tests were performed in all experimental groups as these examinations are considered the proxy of the retrieval stage in the memory formation. It is thought that the retrieval process is influenced by the memory trace intensity as the intensity of experience during the acquisition process in a training session could affect the retrieval memory (1). Thus retrieval stage during the formation of the memory is primarily defined as memory processes, which are requisite for the employment of information already in passive storage, and the retrieval stage is the only part of the memory creation that could be tracked by behavioral studies (14). Hence, a probe trial was utilized for the assessment of the reference memory at the end of learning (15). The probe trial is employed to verify the rats' understanding of the location of the platform and observe the strategy that each rat follows when it discovers the platform is not there (16). We also used the probe trials for the evaluation of the rats' ability to retrieve long-term memory and information learned in the hidden platform test. There are several possible explanations for the findings obtained in our study as mentioned below:




Figure 1. The effect of testosterone on spatial learning. A significant increase was shown in the escape latency in training days in the treatment group compared with the control and sham groups (A). The statistical analysis demonstrated a significant increment in the traveled distance in training days in the testosterone-treated group in comparison to the control and sham groups (B). No significant difference was observed in the swimming speed among the groups (C).

First, testosterone acts as a positive modulator for allosteric function of GABA (17) and also causes the activation of the GABA_A receptor through binding to a particular site present in the interface between β and α subunits (18). Therefore, in our study, testosterone could lead to an increase in the concentration of GABA, as well as impairment in spatial learning.

Second, it has been alleged that memory fixation is empowered by the repetition of particular experience or tasks (19). Thus increasing the frequency of learning repetitions is capable of enhancing the memory fixation. In line with this, an increase in the spacing interval, i.e., the time period between learning repetitions improves long-term memory performance as well (20). Also, in animals that repeatedly encounter a familiar environment, there would be day-to-day turnover in CA1 pyramidal cells representing this environment, which enables the rats to distinguish the representations of different visits (21). It has been shown that behavioral experience alters the neuronal activity, inducing changes in the density of synapses and

dendritic spine (22). Some studies have indicated the increased spine density in CA1 pyramidal cells and basal dendrites in spatially trained rats. Concerning the unchanged length of the dendrites, a higher spine density reflects an expanded number of excitatory synapses per neuron, correlated with spatial learning, as well as altered connectivity. Furthermore, the trained animals showed increased learning ability as shown by faster acquisition in a water maze task. Studies demonstrated that behavioral training could cause a structural change in the hippocampal cortex of rats (23). Dendritic spines play a significant role in the interplay between experience and memory (24). The blockade of synaptic transmission in mature hippocampal slices leads to homeostatic spinogenesis that considerably elevates the number of stubby spines and non-synaptic filopodia, signifying a recapitulation of the early development (25). Therefore, regarding our results, the blockade of synaptic transmission by testosterone might increase the rate of spinogenesis.

Third, it has been demonstrated that LTP is comprised




Figure 2. Average escape latency (A), traveled distance (B), and swimming speed (C) across all training days in the testosterone-treated group. Here, ** $P < 0.01$, *** $P < 0.001$ show significant differences among the experimental groups (A and B). No significant difference was observed among the studied groups considering the swimming speed (C).

of two-time phases in comparison to memory. The short-lasting LTP triggers the long-lasting LTP when multiple trains of high-frequency stimulations are applied (26). Upon the induction of LTP, the number and activity of AMPARs, mediating synaptic transmission, as well as the number of spines would be increased (27). There is a correlation between LTP and the physical enlargement of spines, forming new spines from the neuronal shaft. Hence, the induction of LTP could result in an increase in the density of spines in neurons (28). Therefore, LTP is capable of causing structural changes in synapses (6). After the trigger of LTP, all of the local dendrites, nuclear transcription, and somatic translation incorporate to synthesize proteins required for the maintenance of the functional and structural plasticity. Actually, the occurrence of the LTP expression and maintenance does not sequentially happen;

rather the structural changes such as the increase of spine growth and post-synaptic density would be evident after the induction. Though the early-mentioned structural alterations in spines have been studied in some details, the presence of LTPs may not be necessary as the synapses of the dendrites are capable of expressing and maintaining LTP (29). Studies indicate that LTP is an essential mediator of the induction process in the postsynaptic influx of calcium ions through the activated receptor of NMDA. Since the excitatory synapses of hippocampal pyramidal cells are mainly located on dendritic spines, calcium ions are presumably concentrated on the spine heads (29). In fact, spines could be considered a principle compartment for calcium ion (30).

The last reason is that testosterone is vital for the maintaining of the normal spine synapse density in the CA1




Figure 3. The effect of testosterone on probe test and visible platform test. No significant difference was shown between the treatment group and other groups regarding the time spent in the target quadrant (A) and the distance to the platform (B). There was no significant difference among all groups of the study considering the escape latency (C) or velocity (D) in the visible platform test.

region of the male rats. Testosterone has a considerable impact on the spine synapse density in the CA1; however, this effect is primarily mediated via androgens instead of the estrogen receptors (13). The depletion of testosterone through castration cause a significant reduction in dendritic synapses and the administration of testosterone to castrated rats can retrieve the synaptic density to the baseline levels (31). According to our study, it is hypothesized that testosterone is able to increase dendritic spines and spatial memory.

4.1. Conclusions

Although testosterone causes the perturbations in learning, an increase in the frequency of spatial learning in MWM could lead to the repeated LTP, spinogenesis, increased spine density, and spontaneous generation of new spines, resulting in the improvement in spatial memory. On the other hand, the effect of testosterone through the

androgen receptors causes an increase in dendritic spines leading to improved spatial memory in retrieval test.

Footnotes

Authors' Contribution: Study concept and design: Nasser Naghdi; acquisition of the data: Azadeh Gholaminejad; analysis and interpretation of the data: Azadeh Gholaminejad; drafting of the manuscript: Azadeh Gholaminejad; critical revision of the manuscript for important intellectual content: Mohammad Nasehi; statistical analysis: Azadeh Gholaminejad; administrative, technical, and material support: Hamid Gholamipour-Badie; study supervision: Nasser Naghdi, Hamid Gholamipour-Badie and Mohammad Nasehi.

Conflict of Interests: The authors declare they have no conflict of interests.

Ethical Considerations: ICSS is on the verge of receiving the code of ethics.

Funding/Support: This study was supported by Azadeh Gholaminejad and in partial by Pasteur Institute of Iran.

References

- Quillfeldt JA. Behavioral methods to study learning and memory in rats. In: Andersen M, Tufik S, editors. *Rodent model as tools in ethical biomedical research*. Cham: Springer, Cham; 2016. doi: [10.1007/978-3-319-11578-8_17](https://doi.org/10.1007/978-3-319-11578-8_17).
- Vorhees CV, Williams MT. Assessing spatial learning and memory in rodents. *ILAR J*. 2014;**55**(2):310–32. doi: [10.1093/ilar/ilu013](https://doi.org/10.1093/ilar/ilu013). [PubMed: [25225309](https://pubmed.ncbi.nlm.nih.gov/25225309/)]. [PubMed Central: [PMC4240437](https://pubmed.ncbi.nlm.nih.gov/PMC4240437/)].
- Oe Y, Tominaga-Yoshino K, Hasegawa S, Ogura A. Dendritic spine dynamics in synaptogenesis after repeated LTP inductions: Dependence on pre-existing spine density. *Sci Rep*. 2013;**3**:1957. doi: [10.1038/srep01957](https://doi.org/10.1038/srep01957). [PubMed: [23739837](https://pubmed.ncbi.nlm.nih.gov/23739837/)]. [PubMed Central: [PMC3674431](https://pubmed.ncbi.nlm.nih.gov/PMC3674431/)].
- Shahzad P, Nasser N. GABA_B receptor antagonist (CGP35348) improves testosterone induced spatial acquisition impairment in adult male rat. *J Behav Brain Sci*. 2015;**5**(11):491. doi: [10.4236/jbbs.2015.511047](https://doi.org/10.4236/jbbs.2015.511047).
- Marshad RA, Khatib RA, Amer H, Shammari MA, Otaibi AA, Otaibi FA, et al. Streptozotocin-induced diabetes mellitus affects the NMDA receptors: Role of caffeine administration in enhancing learning, memory and locomotor deficits. *Int J Health Sci (Qassim)*. 2018;**12**(3):10–7. [PubMed: [29896066](https://pubmed.ncbi.nlm.nih.gov/29896066/)]. [PubMed Central: [PMC5969783](https://pubmed.ncbi.nlm.nih.gov/PMC5969783/)].
- Gasbarri A, Pompili A. Involvement of glutamate in learning and memory. In: Meneses A, editor. *Identification of neural markers accompanying memory*. Elsevier; 2014. p. 63–77. doi: [10.1016/B978-0-12-408139-0.00004-3](https://doi.org/10.1016/B978-0-12-408139-0.00004-3).
- Akkerman S, Blokland A, Prickaerts J. Possible overlapping time frames of acquisition and consolidation phases in object memory processes: A pharmacological approach. *Learn Mem*. 2016;**23**(1):29–37. doi: [10.1101/lm.040162.115](https://doi.org/10.1101/lm.040162.115). [PubMed: [26670184](https://pubmed.ncbi.nlm.nih.gov/26670184/)]. [PubMed Central: [PMC4749836](https://pubmed.ncbi.nlm.nih.gov/PMC4749836/)].
- Szapiro G, Galante JM, Barros DM, Levi de Stein M, Vianna MR, Izquierdo LA, et al. Molecular mechanisms of memory retrieval. *Neurochem Res*. 2002;**27**(11):1491–8. doi: [10.1023/A:1021648405461](https://doi.org/10.1023/A:1021648405461). [PubMed: [12512953](https://pubmed.ncbi.nlm.nih.gov/12512953/)].
- Barnhart CD, Yang D, Lein PJ. Using the Morris water maze to assess spatial learning and memory in weanling mice. *PLoS One*. 2015;**10**(4). e0124521. doi: [10.1371/journal.pone.0124521](https://doi.org/10.1371/journal.pone.0124521). [PubMed: [25886563](https://pubmed.ncbi.nlm.nih.gov/25886563/)]. [PubMed Central: [PMC4401674](https://pubmed.ncbi.nlm.nih.gov/PMC4401674/)].
- Martin SJ, Grimwood PD, Morris RG. Synaptic plasticity and memory: An evaluation of the hypothesis. *Annu Rev Neurosci*. 2000;**23**:649–711. doi: [10.1146/annurev.neuro.23.1.649](https://doi.org/10.1146/annurev.neuro.23.1.649). [PubMed: [10845078](https://pubmed.ncbi.nlm.nih.gov/10845078/)].
- McCann RF, Ross DA. A fragile balance: Dendritic spines, learning, and memory. *Biol Psychiatry*. 2017;**82**(2):e11–3. doi: [10.1016/j.biopsych.2017.05.020](https://doi.org/10.1016/j.biopsych.2017.05.020). [PubMed: [28645359](https://pubmed.ncbi.nlm.nih.gov/28645359/)]. [PubMed Central: [PMC5712843](https://pubmed.ncbi.nlm.nih.gov/PMC5712843/)].
- Mahmoud RR, Sase S, Aher YD, Sase A, Groger M, Mokhtar M, et al. Spatial and working memory is linked to spine density and mushroom spines. *PLoS One*. 2015;**10**(10). e0139739. doi: [10.1371/journal.pone.0139739](https://doi.org/10.1371/journal.pone.0139739). [PubMed: [26469788](https://pubmed.ncbi.nlm.nih.gov/26469788/)]. [PubMed Central: [PMC4607435](https://pubmed.ncbi.nlm.nih.gov/PMC4607435/)].
- Leranth C, Petnehazy O, MacLusky NJ. Gonadal hormones affect spine synaptic density in the CA1 hippocampal subfield of male rats. *J Neurosci*. 2003;**23**(5):1588–92. doi: [10.1523/JNEUROSCI.23-05-01588.2003](https://doi.org/10.1523/JNEUROSCI.23-05-01588.2003). [PubMed: [12629162](https://pubmed.ncbi.nlm.nih.gov/12629162/)].
- Dudai Y. The neurobiology of consolidations, or, how stable is the engram? *Annu Rev Psychol*. 2004;**55**:51–86. doi: [10.1146/annurev.psych.55.090902.142050](https://doi.org/10.1146/annurev.psych.55.090902.142050). [PubMed: [14744210](https://pubmed.ncbi.nlm.nih.gov/14744210/)].
- Mendez-Couz M, Conejo NM, Gonzalez-Pardo H, Arias JL. Functional interactions between dentate gyrus, striatum and anterior thalamic nuclei on spatial memory retrieval. *Brain Res*. 2015;**1605**:59–69. doi: [10.1016/j.brainres.2015.02.005](https://doi.org/10.1016/j.brainres.2015.02.005). [PubMed: [25680583](https://pubmed.ncbi.nlm.nih.gov/25680583/)].
- Nunez J. Morris water maze experiment. *J Vis Exp*. 2008;**19**. doi: [10.3791/897](https://doi.org/10.3791/897). [PubMed: [19066539](https://pubmed.ncbi.nlm.nih.gov/19066539/)].
- Carver CM, Reddy DS. Neurosteroid interactions with synaptic and extrasynaptic GABA_A receptors: Regulation of subunit plasticity, phasic and tonic inhibition, and neuronal network excitability. *Psychopharmacology*. 2013;**230**(2):151–88. doi: [10.1007/s00213-013-3276-5](https://doi.org/10.1007/s00213-013-3276-5).
- Sieghart W, Ramerstorfer J, Sarto-Jackson I, Varagic Z, Ernst M. A novel GABA(A) receptor pharmacology: Drugs interacting with the alpha(+) beta(-) interface. *Br J Pharmacol*. 2012;**166**(2):476–85. doi: [10.1111/j.1476-5381.2011.01779.x](https://doi.org/10.1111/j.1476-5381.2011.01779.x). [PubMed: [22074382](https://pubmed.ncbi.nlm.nih.gov/22074382/)]. [PubMed Central: [PMC3417481](https://pubmed.ncbi.nlm.nih.gov/PMC3417481/)].
- Hasegawa S, Sakuragi S, Tominaga-Yoshino K, Ogura A. Dendritic spine dynamics leading to spine elimination after repeated inductions of LTD. *Sci Rep*. 2015;**5**:7707. doi: [10.1038/srep07707](https://doi.org/10.1038/srep07707). [PubMed: [25573377](https://pubmed.ncbi.nlm.nih.gov/25573377/)]. [PubMed Central: [PMC4648349](https://pubmed.ncbi.nlm.nih.gov/PMC4648349/)].
- Kornmeier J, Sosic-Vasic Z. Parallels between spacing effects during behavioral and cellular learning. *Front Hum Neurosci*. 2012;**6**:203. doi: [10.3389/fnhum.2012.00203](https://doi.org/10.3389/fnhum.2012.00203). [PubMed: [22783181](https://pubmed.ncbi.nlm.nih.gov/22783181/)]. [PubMed Central: [PMC3390592](https://pubmed.ncbi.nlm.nih.gov/PMC3390592/)].
- Attardo A, Lu J, Kawashima T, Okuno H, Fitzgerald JE, Bito H, et al. Long-term consolidation of ensemble neural plasticity patterns in hippocampal area CA1. *Cell Rep*. 2018;**25**(3):640–650 e2. doi: [10.1016/j.celrep.2018.09.064](https://doi.org/10.1016/j.celrep.2018.09.064). [PubMed: [30332644](https://pubmed.ncbi.nlm.nih.gov/30332644/)].
- Jasinska M, Siucinska E, Jasek E, Litwin JA, Pyza E, Kossut M. Effect of associative learning on memory spine formation in mouse barrel cortex. *Neural Plast*. 2016;**2016**:9828517. doi: [10.1155/2016/9828517](https://doi.org/10.1155/2016/9828517). [PubMed: [26819780](https://pubmed.ncbi.nlm.nih.gov/26819780/)]. [PubMed Central: [PMC4706958](https://pubmed.ncbi.nlm.nih.gov/PMC4706958/)].
- Moser MB, Trommald M, Andersen P. An increase in dendritic spine density on hippocampal CA1 pyramidal cells following spatial learning in adult rats suggests the formation of new synapses. *Proc Natl Acad Sci U S A*. 1994;**91**(26):12673–5. doi: [10.1073/pnas.91.26.12673](https://doi.org/10.1073/pnas.91.26.12673). [PubMed: [7809099](https://pubmed.ncbi.nlm.nih.gov/7809099/)]. [PubMed Central: [PMC45501](https://pubmed.ncbi.nlm.nih.gov/PMC45501/)].
- Yang G, Lai CS, Cichon J, Ma L, Li W, Gan WB. Sleep promotes branch-specific formation of dendritic spines after learning. *Science*. 2014;**344**(6188):1173–8. doi: [10.1126/science.1249098](https://doi.org/10.1126/science.1249098). [PubMed: [24904169](https://pubmed.ncbi.nlm.nih.gov/24904169/)]. [PubMed Central: [PMC4447313](https://pubmed.ncbi.nlm.nih.gov/PMC4447313/)].
- Bourne J, Harris KM. Do thin spines learn to be mushroom spines that remember? *Curr Opin Neurobiol*. 2007;**17**(3):381–6. doi: [10.1016/j.conb.2007.04.009](https://doi.org/10.1016/j.conb.2007.04.009). [PubMed: [17498943](https://pubmed.ncbi.nlm.nih.gov/17498943/)].
- Villers A, Godaux E, Ris L. Long-lasting LTP requires neither repeated trains for its induction nor protein synthesis for its development. *PLoS One*. 2012;**7**(7). e40823. doi: [10.1371/journal.pone.0040823](https://doi.org/10.1371/journal.pone.0040823). [PubMed: [22792408](https://pubmed.ncbi.nlm.nih.gov/22792408/)]. [PubMed Central: [PMC3394721](https://pubmed.ncbi.nlm.nih.gov/PMC3394721/)].
- Yang Y, Wang XB, Frerking M, Zhou Q. Spine expansion and stabilization associated with long-term potentiation. *J Neurosci*. 2008;**28**(22):5740–51. doi: [10.1523/JNEUROSCI.3998-07.2008](https://doi.org/10.1523/JNEUROSCI.3998-07.2008). [PubMed: [18509035](https://pubmed.ncbi.nlm.nih.gov/18509035/)]. [PubMed Central: [PMC2561912](https://pubmed.ncbi.nlm.nih.gov/PMC2561912/)].
- Sala C. Molecular regulation of dendritic spine shape and function. *Neurosignals*. 2002;**11**(4):213–23. doi: [10.1159/000065433](https://doi.org/10.1159/000065433). [PubMed: [12393947](https://pubmed.ncbi.nlm.nih.gov/12393947/)].
- Luscher C, Malenka RC. NMDA receptor-dependent long-term potentiation and long-term depression (LTP/LTD). *Cold Spring Harb Perspect Biol*. 2012;**4**(6). doi: [10.1101/cshperspect.a005710](https://doi.org/10.1101/cshperspect.a005710). [PubMed: [22510460](https://pubmed.ncbi.nlm.nih.gov/22510460/)]. [PubMed Central: [PMC3367554](https://pubmed.ncbi.nlm.nih.gov/PMC3367554/)].
- Holcman D, Schuss Z, Korkotian E. Calcium dynamics in dendritic spines and spine motility. *Biophys J*. 2004;**87**(1):81–91. doi: [10.1529/biophysj.103.035972](https://doi.org/10.1529/biophysj.103.035972). [PubMed: [15240447](https://pubmed.ncbi.nlm.nih.gov/15240447/)]. [PubMed Central: [PMC1304398](https://pubmed.ncbi.nlm.nih.gov/PMC1304398/)].
- Li M, Masugi-Tokita M, Takanami K, Yamada S, Kawata M. Testosterone has sublayer-specific effects on dendritic spine maturation mediated by BDNF and PSD-95 in pyramidal neurons in the hippocampus CA1 area. *Brain Res*. 2012;**1484**:76–84. doi: [10.1016/j.brainres.2012.09.028](https://doi.org/10.1016/j.brainres.2012.09.028). [PubMed: [23010313](https://pubmed.ncbi.nlm.nih.gov/23010313/)].